

Original Research Article

Alternative Medicines for HIV/AIDS in Resource-Poor Settings: Insight from Traditional Medicines Use in Sub-Saharan Africa

Alfred Maroyi

Medicinal Plants and Economic Development (MPED) Research Centre; Department of Botany, University of Fort Hare, Private Bag X1314, Alice 5700, South Africa

*For correspondence: **Email:** amaroyi@ufh.ac.za; **Tel:** +27406022320, **Fax:** +27866177642

Received: 26 February 2014

Revised accepted: 19 July 2014

Abstract

Purpose: To document the utilization of traditional medicines in managing human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS) opportunistic infections in sub-Saharan Africa.

Methods: This study is based on a review of literature published in scientific journals, books, reports from national, regional and international organizations, theses and conference papers obtained from libraries and electronic search of Google Scholar, ISI Web of Science, MEDLINE, Pubmed, Scopus and Science Direct.

Results: A total of 79 medical conditions related to HIV/AIDS were treated using 74 plant species. The common diseases treated by herbal remedies were bacterial/fungal infections, boosting of appetite/immunity, cold/cough, cryptococcal meningitis, diarrhea, fever, herpes simplex/zoster, oral/oesopharyngeal candidiasis, skin infections/rash, tuberculosis and wounds. More than three-quarters of the documented plant species (63 species, 85.1 %) have anti-HIV active compounds.

Conclusion: This study reveals that traditional medicines are often used as alternative sources of medicines for HIV/AIDS opportunistic infections in sub-Saharan Africa. Further investigations are needed to explore the bioactive compounds of these herbal medicines, aimed at exploring the bioactive compounds that can be developed into anti-HIV drugs.

Keywords: Antiretroviral, HIV/AIDS, Sub-Saharan Africa, Traditional medicines

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

The acquired immunodeficiency syndrome (AIDS), caused by infection with the human immunodeficiency virus (HIV) has become a devastating epidemic in sub-Saharan Africa [1]. Ethnobotanical studies conducted in the following sub-Saharan African countries indicated that traditional healers and local communities extensively use medicinal plants to manage the effects of HIV/AIDS in Cameroon [2], Namibia

[3,4], Rwanda [5], South Africa [6-8], Sudan [9], Tanzania [1,10] and Uganda [11-13].

Despite the increasing acceptance of traditional medicines as an additional and alternative therapeutic strategy in the fight against HIV/AIDS opportunistic diseases in sub-Saharan Africa [1-4,7,8,10-13], this rich indigenous knowledge on traditional remedies is not adequately documented. Documentation of plants used for treating and managing HIV/AIDS opportunistic diseases in sub-Saharan Africa is urgent as there

is need to search for novel anti-HIV compounds. Moreover, an estimated 25 % of pharmaceutical drugs and 11 % of drugs considered essential by WHO are derived from plants; a large number of synthetic drugs obtained from precursor compounds originating from plants [14].

Medicinal plants used for treating and managing HIV/AIDS opportunistic diseases in sub-Saharan Africa are an integral part of a culturally accepted and holistic health care system that has been in existence for several generations. Therefore, traditional medicines are possible sources of complementary and alternative medicines for HIV/AIDS opportunistic diseases.

This review briefly examines the role of herbal medicines in the fight against HIV/AIDS opportunistic diseases in sub-Saharan Africa. It is hoped that this review will provide important and relevant information that will form a basis for further research to identify and isolate bioactive constituents that can be developed as drugs for the treatment and management of HIV/AIDS opportunistic diseases.

EXPERIMENTAL

In this study, the focus was on herbal plants that are used to treat or manage HIV/AIDS opportunistic diseases in sub-Saharan Africa, and also reported to have anti-viral bioactivity based on *in vitro* or *in vivo* studies. Only papers that included information on plant identity, the HIV/AIDS opportunistic diseases treated and plant part(s) used were included in this review.

The International Plant Name Index (www.ipni.org) and the Royal Botanic Garden and Missouri Botanic Garden plant name database (www.theplantlist.org) were used to validate plant scientific names, plant families and plant authorities. In addition to this, the anti-viral efficacy of the documented plant species, either *in vitro* or *in vivo* was also verified from published literature. Therefore, this review draws heavily on the results of ethnobotanical surveys undertaken in contemporary sub-Saharan Africa on traditional medicines used to treat or manage HIV/AIDS opportunistic diseases [1-13], as well as secondary data collected through a review of 60 research reports, policy documents, reports from national, regional and international organizations, and conference papers published in scientific journals (53), books (4), theses (2) and one as an abstract.

Literature was searched on international online databases such as ISI Web of Science, MEDLINE, Science Direct, Scopus and Google Scholar using specific search terms such as “herbal medicines used to treat HIV/AIDS opportunistic diseases”, “medicinal plants used to treat HIV/AIDS opportunistic diseases”, “traditional medicines used to treat HIV/AIDS opportunistic diseases”, and “anti-viral activity of [each documented plant species]”. References were also identified by searching the library collections of the National Herbarium and Botanic Gardens, Zimbabwe and University of Fort Hare, South Africa.

FINDINGS

This study recorded 74 plant species widely used traditionally in treating and managing HIV/AIDS opportunistic diseases in sub-Saharan Africa (Table 1). These medicinal plants were distributed among 37 families and 65 genera. The majority of medicinal plants (50, 67.6 %) used to treat and manage HIV/AIDS opportunistic diseases in sub-Saharan Africa were from 13 families (Table 2). Plant families with the highest number of medicinal plants were Fabaceae *sensu lato* (11 species), followed by Asteraceae and Euphorbiaceae (5 species each), Combretaceae, Lamiaceae and Myrtaceae (4 species each), Amaryllidaceae, Cucurbitaceae and Phyllanthaceae (3 species each), Apiaceae, Celastraceae, Clusiaceae and Xanthorrhoeaceae (2 species each). The rest of the families were represented by one species each (Table 1). The genera with highest number of species were Combretum with three species, followed by *Artemisia*, *Cucurbita*, *Euphorbia*, *Garcinia*, *Plectranthus*, *Senna* and *Syzygium* with 2 species each (Table 1).

Growth form and plant parts used

Trees (43.2 %) were the primary sources of the medicinal plants used for treating and managing HIV/AIDS opportunistic diseases in sub-Saharan Africa, followed by shrubs (27 %), herbs (24.3 %) and climbers (5.4 %) (Table 1). The plant parts used for making herbal preparations were the bark, bulbs, fruits, leaves, leaf sap, roots, stem, tubers and twigs (Table 1). The leaves (67.6 %) were the most frequently used, followed by roots (35.1 %), bark (31.1 %), bulb and seeds (5.4 % each), fruits, stem and twigs (2.7 % each), tubers and leaf sap (1.4 % each) (Table 1).

Table 1: Plant species used to treat and manage various HIV/AIDS opportunistic diseases in sub-Saharan Africa

Family, species name	Growth habit	Part(s) used	Condition treated, country and references in brackets	Pharmacological activities
Amaryllidaceae				
<i>Allium sativum</i> L.	Herb	Bulb	Aspergillosis, oesopharyngeal candidiasis (South Africa) [8], skin infections (Cameroon) [2]	HIV-1 reverse transcriptase inhibitory effect [15]
<i>Crinum macowani</i> Baker	Herb	Bulb	Acne, backache, blood cleansing, boils, cold, fever, glandular swelling, kidney and bladder diseases, rheumatism, wounds, tonic, tuberculosis, venereal diseases (South Africa) [8,16,17]	Anti-HIV activity [16]
<i>Tulbaghia violacea</i> Harv.	Herb	Leaves	Arthritis, asthma, cancer, cold, cough, cryptococcal meningitis, fever, intestinal worms, rheumatism, tuberculosis (South Africa) [8,16,17]	Anti-HIV activity [16]
Anacardiaceae				
<i>Mangifera indica</i> L.	Tree	Bark/leaves	Cough, diarrhea, tuberculosis (Tanzania, Uganda) [1,11-13]	Inhibits HSV-1 and 2 replication [18]
Apiaceae				
<i>Alepidea amatymbica</i> Eckl. & Zeyh.	Shrub	Bark/roots	Aspergillosis, boost appetite, cryptococcal meningitis, diarrhea, clean blood vessels, oesopharyngeal candidiasis (South Africa) [7,8]	Antiretroviral activity [19]
<i>Daucus carota</i> L.	Herb	Leaves	Skin infections (Cameroon) [2]	Inhibits HSV-1 replication [20]
Apocynaceae				
<i>Cascabela thevetia</i> (L.) Lippold	Shrub	Roots	Cough, tuberculosis (Uganda) [11]	HIV-I reverse transcriptase and HIV-I intrase inhibitor activities [21]
Asteraceae				
<i>Artemisia afra</i> Jacq. ex Wild	Shrub	Leaves	Oesopharyngeal candidiasis (South Africa) [8]	Anti-HIV activity [22]
<i>Artemisia annua</i> L.	Shrub	Leaves	Bacterial/fungal infections (Uganda) [11], skin infections (Cameroon) [2]	Anti-HIV activity [22]
<i>Aspilia pluriseta</i> Schweinf.	Herb	Leaves	Kwashiorkor, worms, wounds (Rwanda) [5]	Anti-HIV activity [5]
<i>Tithonia diversifolia</i> (Hemsl.) A. Gray	Shrub	Leaves	Ascariasis, diarrhea (Rwanda) [5]	Anti-HIV activity [5]
<i>Vernonia amygdalina</i> Del.	Shrub	Leaves/roots	Ascariasis, hepatitis, malaria (Rwanda) [5], diarrhea, herpes zoster (Tanzania, Uganda) [1,10,11], cryptococcal meningitis, herpes simplex (Cameroon, Tanzania) [1,2,10], skin infections (Cameroon) [2], boost appetite/immunity, stomach ache, fever (Cameroon, Uganda) [2,11,13], skin rash (Cameroon, Tanzania, Uganda) [1,2,10,13], backache, cough, headache (Cameroon, Uganda) [2,13], stomachache (Uganda) [13]	Anti-HIV activity [23]
Bignoniaceae				
<i>Kigelia africana</i> (Lam.) Benth.	Tree	Bark/fruits	Herpes simplex (Namibia, Tanzania) [1,3,4], diarrhea (Namibia) [3], skin rash (Uganda) [12]	Inhibits HIV-1 reverse transcriptase [24]
Brassicaceae				
<i>Raphanus sativus</i> L.	Herb	Leaves	Skin infections (Cameroon) [2]	Antiviral activity [25]
Canellaceae				
<i>Warburgia ugandensis</i> Sprague	Tree	Bark/leaves/roots	Bacterial/fungal infection, cough, diarrhea, tuberculosis (Uganda) [11]	Inhibits HIV-1 reverse transcriptase [24]
Caricaceae				
<i>Carica papaya</i> L.	Tree	Leaves/roots/seeds	Cough, tuberculosis (Uganda) [11], oral candidiasis (Tanzania) [1,10]	Anti- HIV-1 activity [26]

Celastraceae				
<i>Elaeodendron transvaalense</i> (Burt Davy) R.H. Archer	Tree	Roots	Cold, dysmenorrhea, fever, skin rash (South Africa) [6]	Inhibits HIV-1 reverse transcriptase [6]
<i>Maytenus senegalensis</i> (Lam.) Exell	Tree	Bark/leaves/roots	Bacterial/fungal infections, cough (Uganda) [11], herpes simplex/zoster, oral candidiasis, skin rash, tuberculosis (Uganda, Tanzania) [1,10-12]	Inhibits HIV-1 replication and protease [24]
Cleomaceae				
<i>Cleome gynandra</i> L.	Herb	Leaves	Oral candidiasis (Tanzania) [1,10]	Anti-HIV activity [27]
Clusiaceae				
<i>Garcinia buchananii</i> Bak.	Tree	Bark/roots	Cryptococcal meningitis, diarrhea, herpes simplex/zoster, skin rash, tuberculosis (Namibia, Tanzania) [1,3,4]	HIV-1 protease inhibitory activity [28]
<i>Garcinia livingstonei</i> T. Anderson	Tree	Bark/roots	Cryptococcal meningitis, diarrhea, herpes simplex/zoster, skin rash, tuberculosis (Namibia) [3]	HIV-1 protease inhibitory activity [28]
Combretaceae				
<i>Combretum hartmannianum</i> Schweinf.	Tree	Bark/leaves/stem	Bacterial infections, febrile, jaundice (Sudan) [9]	Inhibits HIV-1 reverse transcriptase [10]
<i>Combretum molle</i> R. Br. ex G. Don	Tree	Bark/roots	Abdominal pains, convulsions, fever, leprosy, snake bites, worms (South Africa) [6], cough, tuberculosis (Uganda) [11]	Inhibits HIV-1 reverse transcriptase [6]
<i>Combretum paniculatum</i> Vent.	Tree	Leaves	Eye diseases, leprosy (Ethiopia) [29]	Inhibits HIV-1 replication [29]
<i>Terminalia sericea</i> Burch. ex DC.	Tree	Bark	Cryptococcal meningitis (Namibia) [4]	Inhibits HIV-1 reverse transcriptase [6]
Cucurbitaceae				
<i>Cucurbita maxima</i> L.	Climber	Leaves	Skin infections (Cameroon) [2]	Anti-HIV activity [30]
<i>Cucurbita pepo</i> L.	Climber	Leaves/roots	Skin infections (Cameroon) [2]	Inhibits HIV-1 reverse transcriptase [31]
<i>Momordica foetida</i> Schumach.	Herb	Leaves	Cough, diarrhea, fever, skin rash (Uganda) [13]	Anti-HIV activity [32]
Euphorbiaceae				
<i>Croton lechleri</i> Müll. Arg.	Tree	Bark	Anaemia, boost appetite, diarrhea (Namibia) [3]	Antiviral activity [33]
<i>Euphorbia hirta</i> L.	Herb	Leaves/roots	Bacterial/fungal infections, herpes zoster (Uganda) [11], dermatophytoses (South Africa) [8]	Inhibits HIV-1, 2 reverse transcriptase [34]
<i>Euphorbia tirucalli</i> L.	Shrub	Leaves	Cough, tuberculosis (Uganda) [11]	Anti-HIV activity [35]
<i>Jatropha curcas</i> L.	Shrub	Leaves	Herpes zoster (Uganda) [11], oral candidiasis, skin rash (Tanzania) [1,10]	Anti-HIV activity [36]
<i>Ricinus communis</i> L.	Shrub	Roots	Abortifacient, abscesses, antihelmentic, arthritis, asthma, dermatitis, diarrhea, fever, flu, toothache, tuberculosis, wounds (South Africa) [6], cough (Tanzania) [1,10]	Inhibits HIV-1 reverse transcriptase [6]
Fabaceae sensu lato				
<i>Abrus precatorius</i> L.	Climber	Bark/leaves/roots	Oral candidiasis (Namibia, Tanzania) [1,3,10]	Inhibits HIV-1 reverse transcriptase [37]
<i>Albizia amara</i> (Roxb.) B. Boiv.	Tree	Leaves	Stomach ache (Namibia) [3]	Inhibits HIV-1 reverse transcriptase [38]
<i>Cajanus cajan</i> (L.) Millsp.	Herb	Stem	Boost appetite/immunity (Uganda) [11], oral candidiasis (Tanzania) [1,10]	Antiviral activity [39]
<i>Dichrostachys cinerea</i> (L.) Wight & Arn	Shrub	Leaves	Oral candidiasis (Namibia) [3]	Inhibits HIV-1 reverse transcriptase [16]
<i>Erythrina abyssinica</i> DC.	Tree	Bark/root	Cough, tuberculosis (Tanzania) [1,10], diarrhea, herpes zoster (Uganda) [11]	Inhibits HIV-1 replication [24]
<i>Peltophorum africanum</i> Sond.	Tree	Bark/roots	Abdominal pains, cough, diarrhea, dysentery, dysmenorrhea, infertility, sore throat, toothache, tuberculosis, wounds (South Africa) [6]	Inhibits HIV-1 reverse transcriptase [6]
<i>Piliostigma thonningii</i> (Schum.) Milne-Redh.	Tree	Bark/roots	Cough (Namibia) [4]	Anti-HIV, HSV-1 and 2 activity [40]

<i>Senna alata</i> (L.) Roxb.	Tree	Leaves	Skin infections (Cameroon) [2]	Inhibits HIV-1 reverse transcriptase [37]
<i>Senna occidentalis</i> (L.) Link	Herb	Leaves/roots	Cough (Namibia) [3,4], diarrhea (Tanzania) [1,10], jaundice, malaria (Sudan) [9]	Inhibits HIV-1 reverse transcriptase [41]
<i>Sutherlandia frutescens</i> R. Br.	Shrub	Leaves/twigs	Backache, cold, chicken pox, diabetes, flu, physical and mental stress, piles, rheumatism, varicose veins (South Africa) [6]	Inhibits HIV-1 reverse transcriptase [6]
Hypoxidaceae				
<i>Hypoxis hemerocallidea</i> (Fisch.) Mey. & Avé-Lall	Herb	Tuber	Cancer, blood cleansing, boost appetite/immunity, oesopharyngeal candidiasis, tuberculosis, vaginal candidiasis, wounds (South Africa) [7,8,17]	Inhibits HIV-1 reverse transcriptase [42]
Lamiaceae				
<i>Leonotis leonurus</i> (L.) R. Br.	Shrub	Leaves/twigs	Asthma, boils, bronchitis, cold, constipation, cough, eczema, flu, hepatitis, high blood pressure, intestinal worms, itching, muscular cramps, skin infections (South Africa) [7,16,17]	Inhibits HIV-1 reverse transcriptase [16]
<i>Ocimum basilicum</i> L.	Herb	Leaves	Skin infections (Cameroon) [2]	Inhibits HIV-1 reverse transcriptase [43]
<i>Plectranthus amboinicus</i> (Lour.) Spreng.	Herb	Leaves	Bacterial/fungal infections, boost appetite/immunity, cough, tuberculosis (Uganda) [11]	Anti-HIV, HSV-1 and 2 activity [44]
<i>Plectranthus barbatus</i> Andr.	Shrub	Leaves	Herpes simplex/zoster, oral candidiasis, tuberculosis (Tanzania) [1,10]	Anti-HIV-1 activity [45]
Malvaceae				
<i>Grewia mollis</i> Juss.	Shrub	Leaves	Diarrhea (Uganda) [11]	Inhibits HIV-1 replication and protease [24]
Meliaceae				
<i>Azadirachta indica</i> A. Juss	Tree	Leaves	Bacterial/fungal infections (Uganda) [11]	Inhibits HIV-1 replication and protease [24]
Melanthaceae				
<i>Bersama abyssinica</i> Boyle	Tree	Bark/leaves/roots	Ascariasis, cholera, diarrhea, dysentery, malaria, rabies, ulcers, worms (Ethiopia) [29]	Inhibits HIV-1 replication [29]
Moringaceae				
<i>Moringa oleifera</i> Lam.	Shrub	Leaves/seeds/	Boost appetite/immunity (Uganda) [11], diarrhea, skin infections, vomiting (Cameroon, Namibia) [2,3]	Anti-HIV activity [46]
Myricaceae				
<i>Morella salicifolia</i> (Hochst. ex A. Rich.) Verdc. & Polhil	Tree	Bark/roots	Boost appetite/immunity (Uganda) [11], cryptococcal meningitis, diarrhea, herpes simplex, tuberculosis (Tanzania) [1,10]	Inhibits HIV-1 replication and protease [24]
Myrsinaceae				
<i>Maesa lanceolata</i> Forssk.	Shrub	Seeds	Bacterial/fungal infections (Uganda) [11]	Inhibits HIV-1 replication [29]
Myrtaceae				
<i>Eucalyptus globulus</i> Labil.	Tree	Leaves	Cough, tuberculosis (Uganda) [11]	Anti-HSV-1 and 2 activity [47]
<i>Psidium guajava</i> L.	Tree	Bark/leaves	Cough, diarrhea (Namibia, Uganda) [3,11-13], tuberculosis (Namibia, Tanzania, Uganda) [1,3,4,11], vaginal candidiasis (South Africa) [8]	HIV-1 protease inhibitory activity [48]
<i>Syzygium cordatum</i> Hochst. ex Krauss	Tree	Bark/leaves	Diarrhea (Namibia) [3], herpes simplex/zoster, skin rash (Tanzania) [1,10]	Anti-HIV activity [49]
<i>Syzygium guineense</i> (Willd) DC.	Tree	Bark	Diarrhea (Namibia, Tanzania) [1,3,10]	Inhibits HIV-1 reverse transcriptase [50]
Olacaceae				
<i>Ximenia americana</i> L.	Shrub	Bark/roots	Contagious diseases, stomach ache, worms (Ethiopia) [29], oral candidiasis (Namibia) [4], skin rash (Tanzania) [1,10]	Inhibits HIV-1 replication [29]
Oleaceae				
<i>Olea europaea</i> L.	Tree	Bulb	Oesopharyngeal candidiasis (South Africa) [8]	Anti-HIV activity [51]
Papaveraceae				
<i>Argemone</i>	Shrub	Leaves/s	Cryptococcal meningitis (Tanzania) [1,10]	Anti-HIV activity [52]

<i>mexicana</i> L.		leaves		
Passifloraceae				
<i>Passiflora edulis</i> Sims	Climber	Leaves	Boost appetite/immunity (Uganda) [11]	Anti-HSV-1 activity [53]
Phyllanthaceae				
<i>Bridelia micrantha</i> Hochst.	Tree	Bark/roots	Abortifacient, diarrhea, sore eyes, stomach ache (South Africa) [6]	Inhibits HIV-1 reverse transcriptase [6]
<i>Flueggea virosa</i> (Robx.Ex.Wild)Vogt	Shrub	Leaves/roots	Cough, tuberculosis (Uganda) [11]	Anti-HIV-1 reverse transcriptase [54]
<i>Phyllanthus reticulatus</i> Poir.	Shrub	Leaves	Herpes simplex (Tanzania) [1,10]	Anti-HIV-1 reverse transcriptase [55]
Polygalaceae				
<i>Securidaca longipedunculata</i> Fres.	Tree	Bark/leaves/roots	Cryptococcal meningitis, oral candidiasis (Namibia, Tanzania) [1,3,10], gonorrhoea (Namibia) [3], cough, syphilis (Ethiopia, Namibia) [3,29], diarrhea, rheumatism, stomach ache, tuberculosis (Ethiopia) [29]	Inhibits HIV replication [56]
Polygonaceae				
<i>Rumex nepalensis</i> Spreng.	Herb	Leaves	Boils, kwashiorkor, worms (Rwanda) [5]	Anti-HIV activity [5]
Rutaceae				
<i>Citrus limon</i> (L.) Burm.f.	Tree	Leaves/fruits/roots	Aspergillosis, cryptococcal meningitis, diabetes, high blood pressure (South Africa) [7,8], tuberculosis (Tanzania) [1,10]	Anti-HIV-1 activity [57]
Sapindaceae				
<i>Dodonaea viscosa</i> (L.) Jacq.	Shrub	Leaves	Cold, fever, malaria, piles, skin infections, sore throat, wounds (Ethiopia) [29]	Inhibits HIV-1 replication [29]
Solanaceae				
<i>Solanum americanum</i> L.	Shrub	Leaves	Boost appetite/immunity (Uganda) [11]	Antiviral activity [58]
Xanthorrhoeaceae				
<i>Aloe vera</i> (L.) Burm.f.	Herb	Leaf sap/leaves	Bacterial/fungal infections, boost appetite/immunity, cough, diarrhea, herpes zoster, tuberculosis (Uganda) [11]	Anti-HSV-2 activity [59]
<i>Bulbine alooides</i> (L.) Willd.	Herb	Leaves/roots	Burns, cracked lips, diarrhea, herpes simplex, itching, skin rash, ringworm, vomiting, wounds (South Africa) [16,17]	Anti-HIV activity [16]
Zingiberaceae				
<i>Zingiber officinalis</i> L.	Herb	Bulb	Aspergillosis (South Africa) [8], boost appetite/immunity, cough, tuberculosis (Uganda) [11], skin infections (Cameroon) [2]	Anti-HIV-1 activity [60]

Table 2: Families with the largest number of medicinal plants (more than 2 species) used to treat and manage HIV/AIDS opportunistic diseases in sub-Saharan Africa

Family	Number of medicinal plants	%
Fabaceae <i>sensu lato</i>	11	14.9
Asteraceae	5	6.8
Combretaceae	4	5.4
Lamiaceae	4	5.4
Myrtaceae	4	5.4
Amaryllidaceae	3	4.1
Cucurbitaceae	3	4.1
Phyllanthaceae	3	4.1
Apiaceae	2	2.7
Celastraceae	2	2.7
Clusiaceae	2	2.7
Xanthorrhoeaceae	2	2.7

Table 3: Major ailment and disease categories and plant species reported. Most species were reported in more than one ailment and disease category.

Ailment/disease	Number of species
Wounds	8
Bacterial/fungal infections	9
Fever	9
Boost immunity	11
Boost appetite	13
Cryptococcal meningitis	13
Oral/oesopharyngeal candidiasis	19
Herpes simplex/zoster	27
Tuberculosis	30
Skin infections/rash	31
Diarrhea	32
Cold/cough	33

HIV/AIDS opportunistic diseases and ailments treated with herbal medicines

A total of 79 medical conditions related to HIV/AIDS were treated using herbal remedies (Table 1). Bacterial/fungal infections, boosting of appetite/immunity, cold/cough, cryptococcal meningitis, diarrhea, fever, herpes simplex/zoster, oral/oesopharyngeal candidiasis, skin infections/rash, tuberculosis and wounds were treated with the highest number of medicinal plant species (Table 3).

DISCUSSION

The checklist in Table 1 provides baseline data on medicinal plants that are currently used to treat and manage HIV/AIDS opportunistic diseases in sub-Saharan Africa. This inventory is a crucial starting point in trying to assess the efficacy of traditional medicines in the management of HIV/AIDS opportunistic diseases as well as the development of effective drugs for the treatment of HIV/AIDS. One of the possible approaches to finding novel anti-HIV therapeutic agents is the screening of medicinal plants that are widely used in local communities to treat and manage HIV/AIDS opportunistic diseases.

Out of 74 medicinal plants widely used in sub-Saharan Africa to treat and manage HIV/AIDS opportunistic diseases, 63 species (85.1 %) have proven anti-HIV active compounds and known modes of action (Table 1). Among these is *Kigelia africana* used to treat diarrhea and herpes simplex in Namibia [3], herpes simplex in Tanzania [1] and skin rash in Uganda [12]. *Maytenus senegalensis* is used as herbal medicine for bacterial/fungal infections and cough in Uganda [11], and to treat herpes simplex/zoster, oral candidiasis, skin rash and tuberculosis in both Tanzania and Uganda [1,10-12]. Similarly, *Garcinia burchananii* has the same ethnomedicinal applications in Namibia [3] and Tanzania [1] as herbal medicine for chronic diarrhea, cryptococcal meningitis, herpes simplex/zoster, skin rash and tuberculosis [1,3,10].

Another noteworthy plant species used for treating and managing HIV/AIDS opportunistic diseases in sub-Saharan Africa are *Hypoxis hemerocallidea* and *Sutherlandia frutescens* (Table 1). These two species are currently recommended by the South African Ministry of Health for HIV management [6,17,42]. This is in accordance with the World Health Organization's recommendation that traditional medicines

should be included in national responses to HIV/AIDS pandemic [14]. Some of these species can, therefore, be targeted for phytochemical and pharmacological studies with the aim of identifying active ingredients contained by such plants required for treating and managing HIV/AIDS. Such research is important, as there is need to search for improved antiretroviral agents which can be added to or replace the current drugs in use.

Despite the presence of anti-HIV activity in more than 85 % of the plant species in Table 1, these plant species have not been fully explored. Further investigations on phytochemical constituents and subsequent screening are needed for opening new opportunities to develop pharmaceutical drugs based on anti-HIV constituents shown by some of these plant species.

This study revealed that many people in sub-Saharan Africa use traditional medicines for the treatment and management of HIV/AIDS opportunistic diseases despite the increasing availability of antiretroviral therapy. People in sub-Saharan Africa, particularly the poor are still dependent on medicinal plants for the treatment of basic human ailments like backache, cold, cough, diarrhea, fever, headache, skin infections and wounds [14,17,24,44]. Therefore, traditional medicines remain the main source of primary health care available and accessible to all people in sub-Saharan Africa.

Although no plant-derived drug is currently in clinical use to treat and manage HIV/AIDS opportunistic diseases, knowledge on these plants is very important as this can serve as leads in the discovery of new anti-HIV agents. Medicinal plants may yield novel compounds that may be of interest for HIV/AIDS drug development particularly those medicinal plants characterized by anti-HIV activity (Table 1).

Limitations of the study

This study has taken a focused approach and generated extensive exploratory and qualitative data, however, the following limitations are acknowledged in relation to its findings. Detailed accounts of plants used to treat and manage HIV/AIDS opportunistic diseases are available from Cameroon, Namibia, Rwanda, South Africa, Sudan, Tanzania and Uganda only. Although, the results from these countries are diverse in nature, they may not apply directly to all countries in the sub-Saharan Africa. The

advantages of this limited sample size and sharp focus on detailed accounts on medicinal plants used to treat and manage HIV/AIDS opportunistic diseases in Cameroon, Namibia, Rwanda, South Africa, Sudan, Tanzania and Uganda resulted in a better and deeper understanding of traditional medicines useful for HIV/AIDS management in sub-Saharan Africa. Overall, while the study is useful in gaining an understanding of utilization of traditional medicines in managing HIV/AIDS opportunistic infections in sub-Saharan Africa and the value of traditional medicines in a wide spectrum of countries in the region, it is clear that more detailed national level studies should be undertaken. The present study provided a sample of preliminary data that could open up pertinent questions relating to the management of HIV/AIDS opportunistic infections using herbal medicines.

CONCLUSION

Correlation between the ethnomedicinal employment and the pharmacological activities have been duly observed and described in this study. In fact, this study suggests that some of the plant species have the potential to be developed as anti-HIV agents. While there are still gaps in the phytochemistry and pharmacological studies conducted so far, which need to be bridged in order to exploit the full medicinal potential of these species, it is still very clear that these plant species have tremendous potential for the future. This inventory will assist future workers on the selection of herbal plants to evaluate for phytochemical safety and pharmaceutical efficacy. There is also need for more research on the active compounds of these herbal medicines, some of which have already shown interesting pharmacological activities as shown in Table 1.

REFERENCES

1. Kisangau DP, Lyaruu HV, Hosea KM, Joseph CC. Use of traditional medicines in the management of HIV/AIDS opportunistic infections in Tanzania: a case in the Bukoba rural district. *J Ethnobiol Ethnomed* 2007; 3: 29.
2. Noumi N, Manga PN. Traditional medicines for HIV/AIDS and opportunistic infections in North-West Cameroon: case of skin infections. *Amer J Trop Med Pub Health* 2011; 1(3): 44-64.
3. Chinsembu KC, Hedimbi M. An ethnobotanical survey of plants used to manage HIV/AIDS opportunistic infections in Katima Mulilo, Caprivi region, Namibia. *J Ethnobiol Ethnomed* 2010; 6: 25.
4. Hedimbi M, Chinsembu KC. Ethnomedicinal study of plants used to manage HIV/AIDS-related disease conditions in the Ohangwena region, Namibia. *Int J Med PI Res* 2012; 1(1): 4-11.
5. Cos P, Hermans N, De Bruyne T, Apers S, Sindambiwe JB, Witvrouw M, De Clercq E, Van den Berghe D, Pieters L, Vlietinck AJ. Antiviral activity of Rwandan medicinal plants against human immunodeficiency virus type-1 (HIV-1). *Phytomed* 2002; 9: 62-68.
6. Bessong PO, Obi CL, Andréola M-L, Rojas LB, Pouységu L, Igumbor E, Meyer JJ, Quideau S, Litvak S. Evaluation of selected South African medicinal plants for inhibitory properties against human immunodeficiency virus type 1 reverse transcriptase and integrase. *J Ethnopharmacol* 2005; 99(1): 83-91.
7. Omoruyi BE, Bradley G, Afolayan AJ. Ethnomedicinal survey of medicinal plants used for the management of HIV/AIDS infection among local communities of Nkonkobe Municipality, Eastern Cape, South Africa. *J Med PI Res* 2012; 6(19): 3603-3608.
8. Otang WM, Grierson DS, Ndip R. Ethnobotanical survey of medicinal plants used in the management of opportunistic fungal infections in HIV/AIDS patients in the Amathole District of the Eastern Cape Province, South Africa. *J Med PI Res* 2012; 6(11): 2071-2080.
9. Ali H, Konig GM, Khalid SA, Wright AD, Kaminsky R. Evaluation of selected Sudanese medicinal plants for their *in vitro* activity against hemoflagellates, selected bacteria, HIV-1-RT and tyrosine kinase inhibitory and for cytotoxicity. *J Ethnopharmacol* 2002; 83(3): 219-228.
10. Kisangau DP, Herrmann TM, Lyaruu HVM, Hosea KM, Joseph CC, Mbwambo ZH, Masimba PJ. Traditional knowledge, use practices and conservation of medicinal plants for HIV/AIDS care in rural Tanzania. *Ethnobot Res Appl* 2011; 9: 43-57.
11. Asiiimwe S, Kamatenesi-Mugisha M, Namutebi A, Borg-Karlsson A-K, Musiimenta P. Ethnobotanical study of nutri-medicinal plants used for the management of HIV/AIDS opportunistic ailments among the local communities of western Uganda. *J Ethnopharmacol* 2013; 150(2): 639-648.
12. Lamorde M, Tabuti JR, Obua C, Kukunda-Byobona C, Lanyero H, Byakika-Kibwika P, Bbosa GS, Lubenga A, Oqwal-Okeng J, Ryan M, Waako PJ, Merry C. Medicinal plants used by traditional medicine practitioners for the treatment of HIV/AIDS and related conditions in Uganda. *J Ethnopharmacol* 2010; 130(1): 43-53.
13. Langlois-Klassen D, Kipp W, Jhangri GS, Rubaale T. Use of traditional herbal medicine by AIDS patients in Kabarole District, Western Uganda. *Am J Trop Med Hyg* 2007; 77(4): 757-763.
14. Rates SMK. Plants as a source of drugs. *Toxicon* 2001; 39: 603-613.

15. Silprasit K, Seetaha S, Pongsanarakul P, Hannongbua S, Choowongkamon K. Anti-HIV-1 reverse transcriptase activities of hexane extracts from some Asian medicinal plants. *J Med Pl Res* 2011; 5(17): 4194-4201.
16. Klos M, van de Venter M, Milne PJ, Traore HN, Meyer D, Oosthuizen V. In vitro anti-HIV activity of five selected South African medicinal plant extracts. *J Ethnopharmacol* 2009; 124(2): 182-188.
17. van Wyk BE, van Oudtshoorn B, Gericke N. Medicinal plants of South Africa. Pretoria: Briza Publications; 2000. 336 p.
18. Shah KA, Patel MB, Patel RJ, Parmar PK. *Mangifera Indica* (Mango). *Pharmacogn Rev* 2010; 4(7): 42-48.
19. Louvel S, Moodley N, Seibert I, Steenkamp P, Nthambeleni R, Vidal V, Maharaj V, Klimkait T. Identification of compounds from the plant species *Alepidea amatymbica* active against HIV. *South Afr J Bot* 2013; 86: 9-14
20. Torkey ZA. Antiviral activity of polyphenols extracts from *Daucus carota* against Herpes simplex virus type 1. *TOJSAT: The Online J Sci Technol* 2013; 3(1): 20-32.
21. Tewtrakul S, Nakamura N, Hattori M, Fujiwara T, Supavita T. Flavanone and flavonol glycosides from the leaves of *Thevetia peruviana* and their HIV-1 reverse transcriptase and HIV-1 integrase inhibitory activities. *Chem Pharm Bull (Tokyo)* 2002; 50(5): 630-635.
22. Lubbe A, Seibert I, Klimkait T, van der Kooy F. Ethnopharmacology in overdrive: the remarkable anti-HIV activity of *Artemisia annua*. *J Ethnopharmacol* 2012; 141(3): 854-859.
23. Johnson CE, Lin L-Z, Harnly JM, Oladeinde FO, Kinyua AM, Michelin R, Bronner Y. Identification of the phenolic components of *Vernonia amygdalina* and *Russelia equisetiformis*. *J Nat Prod* 2011; 4: 57-64.
24. Rukanga GM, Kofi-Tsekpo MW, Kurokawa M, Kageyama S, Mungai GM, Muli JM, Tolo FM, Kibaya RM, Muthaura CN, Kanyara JN, Tukei PM, Shiraki K. Evaluation of the HIV-1 reverse transcriptase inhibitory properties of extracts from some medicinal plants in Kenya. *Afr J Health Sci* 2002; 9(1): 81-90.
25. Gutiérrez RMP, Perez RL. *Raphanus sativus* (Radish): Their chemistry and biology. *The Sci World J* 2004; 4: 811-837.
26. Rashed K, Luo M-T, Zhang L-T, Zheng Y-T. Phytochemical screening of the polar extracts of *Carica papaya* Linn. and the evaluation of their anti-HIV-1 activity. *J Appl Industrial Sci* 2013; 1(3): 49-53.
27. Mnzava NA, Chigumira Ngwerume F. *Cleome gynandra* L. In: Grubben GJH, Denton OA (eds). PROTA 2: Vegetables/Légumes. [CD-Rom]. PROTA, Wageningen, Netherlands; 2003.
28. Magadula JJ, Tewtrakul S. Anti-HIV-1 protease activities of crude extracts of some *Garcinia* species growing in Tanzania. *Afr J Biot* 2010; 9(12): 1848-1852.
29. Asres K, Bucar F, Karting T, Witrouw M, Pannecouque C, Clercq ED. Antiviral activity against human immunodeficiency virus type 1 (HIV-1) and type 2 (HIV-2) of ethnobotanically selected Ethiopian medicinal plants. *Phytother Res* 2001; 15(1): 62-69.
30. Daly NL, Koltay A, Gustafson KR, Boyd MR, Casas-Finet JR, Craik DJ. Solution structure by NMR of circulin A: A macrocyclic knotted peptide having anti-HIV activity. *J Mol Biol* 1999; 285(1): 333-345.
31. Gerhäuser C, Samtleben R, Tau GT, Pezzuto JM, Lottspeich F, Wagner H. Peponin, a new ribosome-inactivating protein isolated from seeds of *Curcubita pepo* L. inhibits human immunodeficiency virus type 1 reverse transcriptase. *Pharm Pharmacological Lett* 1993; 3: 71-75.
32. Bourinbaiar AS., Lee-Huang S. Potentiation of anti-HIV activity of anti-inflammatory drugs, dexamethasone and indomethacin, by MAP30, the antiviral agent from bitter melon. *Biochem Biophys Res. Commun* 1995; 208(2): 779-885.
33. Williams JE. Review of Antiviral and immunomodulating properties of plants of the Peruvian rainforest with a particular emphasis on Uña de Gato and Sangre de Grado. *Altern Medicine Rev* 2001; 6(6): 567-579.
34. Gyuris A, Szlávik L, Minárovits J, Vasas A, Molnár J, Hohmann J. Antiviral activities of extracts of *Euphorbia hirta* L. against HIV-1, HIV-2 and SIVmac251. *In Vivo* 2009; 23(3): 429-432.
35. Upadhyay B, Singh KP, Kumar A. Ethno-medicinal, phytochemical and antimicrobial studies of *Euphorbia tirucalli* L. *J Phytology* 2010; 2(4): 65-77.
36. Kaur R, Kharb R. Anti-HIV potential of medicinally important plants. *Int J Pharma Bio Sciences* 2011; 2(3): 387-398.
37. Tewtrakul S, Subhadhirasakul S, Rattanasuwan P. HIV-1 protease inhibitory effects of some selected plants in Caesalpiniaceae and Papilionaceae families. *Songklanakarin J Sci Technol* 2003; 25(4): 509-514.
38. Mar W, Tan GT, Cordell GA, Pezzuto JM, Jurcic K, Offermann F, Redl K, Steinke B, Wagner H. Biological activity of novel macrocyclic alkaloids (budmunchiamines) from *Albizia amara* detected on the basis of interaction with DNA. *J Nat Prod* 1991; 54(6): 1531-1542.
39. Nwodo UU, Ngene AA, Iroegbu CU, Onyedikachi OAL, Chigor VN, Okoh AI. In vivo evaluation of the antiviral activity of *Cajanus cajan* on measles virus. *Arch Virol* 2011; 156: 1551-1557
40. Lemessa, D. *Piliostigma thonningii* (Schumach.) Milne-Redh. In: Brink M, Achigan-Dako EG. (eds). Prota 16: Fibres/Plantes à fibres. [CD-Rom]. PROTA, Wageningen, Netherlands; 2010.
41. Lunavath V, Estari M. Inhibition of human immunodeficiency virus (HIV-1) reverse transcriptase by *Cassia occidentalis* L. plant extract. *Int J Sci Eng Res* 2012; 3(7): 1-4.
42. Klos M. In vitro anti-HIV and immunomodulatory activity of five selected South African medicinal plant extracts [dissertation]. [Port Elizabeth]: Nelson Mandela Metropolitan University; 2006. 184 p.

43. Yamasaki K, Nakano M, Kawahata T, Mori H, Otake T, Ueba N, Oishi I, Inami R, Yamane M, Nakamura M, Murata H, Nakanishi T. Anti-HIV-1 activity of herbs in Labiatae. *Biol Pharm Bull* 1998; 21(8): 829-833.
44. Williams CJ. *Medicinal plants in Australia: An antipodean apothecary*. Rosenberg Publishing, NSW, Australia; 2013
45. Kapewangolo P, Hussein AA, Meyer D. Inhibition of HIV-1 enzymes, antioxidant and anti-inflammatory activities of *Plectranthus barbatus*. *J Ethnopharmacol* 2013; 149(1): 184-190.
46. Nworu C S, Okoye EL, Ezeifeke GO, Esimone CO. Extracts of *Moringa oleifera* Lam. showing inhibitory activity against early steps in the infectivity of HIV-1 lentiviral particles in a viral vector-based screening. *Afr J Biotechnol* 2013; 12(30): 4866-4873.
47. Schnitzler P, Schon K, Reichling J. Antiviral activity of Australian tea tree oil and eucalyptus oil against herpes simplex virus in cell culture. *Pharmazie* 2001; 56: 343-347.
48. Mao QC, Zhou YC, Li RM, Hu YP, Liu SW, Li XJ. Inhibition of HIV-1 mediated cell-cell fusion by saponin fraction from *Psidium guajava* leaf. *Zhong Yao Cai* 2010; 33: 1751-1754.
49. Hurinanthan V. *Anti-HIV activity of selected South African medicinal plants [dissertation]*. Durban University of Technology, Durban; 2013. 169 p.
50. Moll A, Heyman HM, Meyer JJM. Plants with activity against the live HI virus and the enzyme, reverse transcriptase. *South Afr J Bot* 2013; 86: 148.
51. Qian Y, Guan T, Tang X, Huang L, Huang M, Li Y, Sun H. Maslinic acid, a natural triterpenoid compound from *Olea europaea*, protects cortical neurons against oxygen-glucose deprivation-induced injury. *Eur J Pharmacol* 2011; 670(1): 148-153.
52. Chang YC, Hsieh PW, Chang FR, Wu RR, Liaw CC, Lee KH, Wu YC. Two new protopines argemexicaines A and B and the anti-HIV alkaloid 6-acetyldihydrochelerythrine from formosan *Argemone mexicana*. *Planta Med* 2003; 69(2): 148-152.
53. Müller V, Chávez JH, Reginatto FH, Zucolotto SM, Niero R, Navarro D, Yunes RA, Schenkel EP, Barardi CRM, Zanetti CR, Simões CMO. Evaluation of antiviral activity of South American plant extracts against herpes simplex virus type 1 and rabies virus. *Phytother Res* 2007; 21(10): 970-974.
54. Wiwat C, Kwantrairat S. HIV-1 reverse transcriptase inhibitors from Thai medicinal plants and *Elephantopus scaber* Linn. *Mahidol University J Pharm Sci* 2013; 40(3): 35-44.
55. Eldeen IMS, Seow E-M, Abdullah R, Sulaiman SF. In vitro antibacterial, antioxidant, total phenolic contents and anti-HIV-1 reverse transcriptase activities of extracts of seven *Phyllanthus* sp. *South Afr J Bot* 2011; 77(1): 75-79
56. Muazu J, Kaita AH. A review of traditional plants used in the treatment of epilepsy amongst the Hausa/Fulani tribes of northern Nigeria. *Afr J Trad Complement Altern Med* 2008; 5(4): 387-390.
57. Lackman-Smith CS, Snyder BA, Marotte KM, Osterling MC, Mankowski MK, Jones M, Nieves-Duran L, Richardson-Harman N, Cummins JE, Sanders-Beer BE. Safety and anti-HIV assessments of natural vaginal cleansing products in an established topical microbicides in vitro testing algorithm. *AIDS Res Therapy* 2010; 7: 22.
58. Javed T, Ashfaq UA, Riaz S, Rehman S, Riazuddin S. In vitro antiviral activity of *Solanum nigrum* against Hepatitis C Virus. *Virology Journal* 2011; 8:26
59. Zandi K, Zadeh MA, Sartavi K, Rastian Z. 2007. Antiviral activity of *Aloe vera* against herpes simplex virus type 2: An in vitro study. *Afr J Biotechnol* 2007; 6(15): 1770-1773.
60. Feng T, Su J, Ding ZH, Zheng YT, Li Y, Leng Y, Liu JK. Chemical constituents and their bioactivities of "Tongling White Ginger" (*Zingiber officinale*). *J Agric Food Chem* 2011; 59(21): 11690-11695.