

Research Article

Methanol Extract of *Polyopes lancifolius* Suppresses Tumor Necrosis Factor- α -Induced *Matrix* Metalloproteinase-9 Expression in T24 Bladder Carcinoma Cells

R.G.P.T. Jayasooriya¹, Taek-Jeong Nam², Gi-Young Kim^{1*} and Yung Hyun Choi^{3,4*}

¹Laboratory of Immunobiology, Department of Marine Life Sciences, Jeju National University, Jeju 690-756, ²Department of Food Science and Biotechnology, Pukyong National University, Busan 608-737, ³Department of Biochemistry and Research Institute of Oriental Medicine, Dongeui University College of Oriental Medicine, Busan 614-054, ⁴Department of Biomaterial Control (BK21 program) and Blue-Bio Industry Regional Innovation Center, Dongeui University, Busan 614-714, Republic of Korea

Abstract

Purpose: To investigate the effects of the methanol extract of *Polyopes lancifolius* (MEPL) on the expression of matrix metalloproteinase-9 (MMP-9) and invasion in T24 human bladder carcinoma cells.

Methods: Reverse transcriptase-polymerase chain reaction (RT-PCR) and Western blot analyses were performed to assess the expression of MMP-9 and its regulatory proteins. MMP-9 activity was evaluated using zymography while matrigel infiltration was performed to assess T24 bladder carcinoma invasion. Electrophoretic mobility assay was used to investigate the nuclear factor- κ B (NF- κ B) activity.

Results: The expression and activity of MMP-9 were significantly increased in response to TNF- α , but MEPL suppressed TNF- α -induced MMP-9 expression and activity. MEPL also inhibited TNF- α -induced MMP-9 expression at the transcriptional level by blocking the activation of the NF- κ B signaling pathway. Furthermore, the extract suppressed TNF- α -induced phosphorylation of I κ B α and consequently sustained cytosolic p65 and p50 expression. Matrigel invasion assay showed that MEPL significantly reduced TNF- α -induced invasion of T24 bladder carcinoma cells.

Conclusion: Collectively, these data indicate that MEPL regulates TNF- α -induced MMP-9 expression by suppressing NF- κ B activity.

Keywords: *Polyopes lancifolius*, *Matrix metalloproteinase-9*, *Invasion*; *Nuclear factor- κ B*

Received: 12 October 2011

Revised accepted: 17 April 2012

*Corresponding author: **Email:** immunkim@jejunu.ac.kr, choiyh@deu.ac.kr; **Tel:** +82 64 754 3427; **Fax:** +82 64 756 3493

INTRODUCTION

Malignant growth of cells in the urinary bladder tissue leads to bladder cancer. Most complications of bladder cancer are attributed to tumor invasion of distant organs, including the regional lymph nodes, lungs, bones, adrenal glands, and intestines [1]. This invasion occurs in multiple steps: tumor cell proliferation, invasion, angiogenesis, intravasation, survival in the circulation, adhesion to endothelial cells, extravasation, and growth in distant organs [2,3].

Matrix metalloproteinases (MMPs) are zinc-dependent endopeptidases that are collectively capable of degrading all types of extracellular matrix proteins as well as cleaving cell surface receptors, releasing apoptotic ligands, and activating chemokines and cytokines [4,5]. In particular, MMPs are expressed in nearly all tumors, wherein they facilitate tumor growth, invasion, and metastasis [6]. Among them, MMP-9 is particularly known to play a critical role in bladder cancer progression, including the aspects of angiogenesis, tumor growth, invasion, and distant metastasis. Induction of MMP-9 by growth factors, such as the tumor necrosis factor- α (TNF- α), has been reported to contribute to enhanced cell migration and invasion [7]. In contrast, it has been recently shown that several plant extracts have the ability to decrease MMP-9 secretion and suppress the invasiveness of cancer cells [8]. In addition, a recent study showed that the nuclear factor-kappa B (NF- κ B) pathway tightly regulates the expression of MMP-9 in several types of cancer cells [9]. Therefore, NF- κ B is considered as a good target to suppress MMP-9 expression in order to inhibit the invasion and metastasis of human bladder cancer.

Polyopes lancifolius is a type of seaweed with medicinal value, which is usually found in the Republic of Korea and Japan [10,11]. Our previous study established the anti-inflammatory effect of the methanol extract of *P. lancifolius* (MEPL) on lipopolysaccharide

(LPS)-stimulated BV2 microglia cells (unpublished). However, the role of MEPL on the regulation of MMP-9 expression in cancer cells is still unknown. Therefore, in this study, we investigated the ability of MEPL to inhibit MMP-9 expression via suppression of the NF- κ B signaling pathway. We also assessed the mechanism by which MEPL regulates the DNA-binding activity of NF- κ B and its downstream gene, MMP-9.

EXPERIMENTAL

Preparation of MEPL

MEPL was purchased from Jeju HI-Tech Industry Development Institute (extract no. 1340; Jeju, Republic of Korea). The red alga, *P. lancifolius* (stock no. AR038) was collected along the Jeju Island coast of Republic of Korea in April, 2005. Briefly, fresh *P. lancifolius* was washed three times with tap water to remove salt, epiphyte and sand on the surface of the samples before storage -20 °C. The frozen samples were lyophilized and homogenized using a grinder before extraction. The dried powder was extracted with 80 % methanol and evaporated in vacuo.

Reagents

TNF- α and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (MTT) were purchased from Sigma Chemical Co, St Louis, MO, USA. Roswell Park Memorial Institute medium (RPMI) and fetal bovine serum (FBS) were obtained from WelGENE Inc. (Daegu, Republic of Korea). Antibodies against p65, p50 and phospho (p)-I κ B α were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Antibody against β -actin was from Sigma. Peroxidase-labeled goat anti-rabbit immunoglobulin was purchased from KOMA Biotechnology (Seoul, Republic of Korea). Other chemicals used were purchased from Sigma.

Cell culture and sample treatment

Human bladder cancer cell line T24 were

cultured at 37 °C in 5 % CO₂ in RPMI medium supplemented with 10 % FBS and antibiotics. For the analysis of cell viability, the cells (1 × 10⁵ cells/ml) were incubated with the various concentrations of MEPL 1 h before stimulation with TNF-α (20 ng/ml) for 24 h.

Cell viability assay

Cell viability was determined by an MTT assay. In brief, T24 bladder carcinoma cells (1 × 10⁵ cells/ml) were plated onto 24 well plates and incubated overnight. The cells were treated with the indicated concentrations of MEPL for 1 h and then stimulated with TNF-α (20 ng/ml). After 24 h, the cells were incubated with a solution of 0.5 mg/ml MTT for 45 min at 37 °C and 5 % CO₂. The supernatant was removed and the formation of formazan was observed by monitoring the signal at 540 nm using a microplate reader.

Isolation of total RNA and RT-PCR

Total RNA was extracted using easy-BLUE™ total RNA extraction kit (Invitrogen, Carlsbad, USA) according to the manufacturer's instruction. Two microgram RNA was reverse-transcribed using MMLV reverse transcriptase (Promega, Madison, WI). cDNA was amplified by PCR using specific primer MMP-9 (forward 5'-gta ttt gtt caa gga tgg gaa ata c-3' and reverse 5'-gca gga tgt cat agg tca cgt ag-3') and GAPDH (forward 5'-tgt gat ggt ggg aay ggg tc-3' and reverse 5'-ttt gat gtc acg cac gat tt-3'). Reaction products were analyzed on 1.0 % agarose gels, and the bands were visualized by ethidium bromide.

Gelatin substrate gel zymography

The cells were incubated at 37 °C in 5 % CO₂ in serum free RPMI medium supplemented with 10 % FBS and antibiotics in the absence or presence of MEPL for 24 h. The supernatants were collected and then subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-

PAGE) in 10 % polyacrylamide gels copolymerized with 1 mg/ml of gelatin. After electrophoresis, the gels were washed several times in 2.5 % Triton X-100 for 1 h at room temperature to remove SDS and then incubated for 24 h at 37 °C in reaction buffer containing 5mM CaCl₂ and 1μM ZnCl₂. The gels were stained with Coomassie blue (0.25 %) for 30 min and then destained for 1 h in a solution of acetic acid and methanol. The proteolytic activity was evidenced as clear bands (zones of gelatin degradation) against the blue background of stained gelatin.

Matrigel invasion assay

Invasion assays were performed using Boyden chambers with polycarbonate nucleopore membrane (Corning, Corning, NY, USA). Cells were trypsinized and 5 × 10⁴ cells were placed onto matrigel-coated transwell for 3 h. The cells were treated with 100 μg/ml of MEPL for 1 h and then stimulated with TNF-α (20 ng/ml). After incubation for 24 h at 37 °C in 5 % CO₂, noninvasive cells in upper chamber were removed with a cotton swab. Invaded cells on the lower surface of the filter were fixed and stained with 0.125 % Coomassie blue. Random fields were counted under a light microscope.

Western blot analysis

Total cell extracts were prepared using PRO-PREP protein extraction kit (iNtRON Biotechnology; Sungnam, Republic of Korea). Briefly, after treatment with the indicated concentrations of MEPL, cells were harvested, washed once with ice-cold PBS and gently lysed for 15 min in 100 μl ice-cold PRO-PREP lysis buffer. Lysates were centrifuged at 14,000 g at 4 °C for 10 min to obtain the supernatants. Supernatants were collected and protein concentrations were determined using a Bio-Rad protein assay kit (Bio-Rad, Hercules, CA, USA). The proteins were separated on SDS-polyacrylamide gels, transferred to nitrocellulose membranes (Schleicher & Schuell, Keene, NH), and detected using an enhanced chemilu-

minescence detection system (model RPN2108, Amersham, Arlington Heights, IL, USA).

Electrophoretic mobility assay (EMSA)

DNA-protein binding assays were carried out with nuclear extract. Synthetic complementary NF- κ B (5'-agt tga ggg gac ttt ccc agg c-3') binding oligonucleotides (Santa Cruz Biotechnology) were 3'-biotinylated using the biotin 3'-end DNA labeling kit (Pierce) according to the manufacturer's instructions, and annealed for 30 min at room temperature. Assays were loaded onto native 4 % polyacrylamide gels pre-electrophoresed for 60 min in 0.5 \times Tris borate/EDTA before being transferred onto a positively charged nylon membrane (HybondTM-N+) in 0.5 \times Tris borate/EDTA at 100 V for 30 min. The transferred DNAs were cross-linked to the membrane at 120 mJ/cm². Horseradish peroxidase-conjugated streptavidin was used according to the manufacturer's instructions to detect the transferred DNA.

Statistical analysis

All data were derived from at least three independent experiments. Statistical analyses were conducted using SigmaPlot software (version 11.0). Values were presented as mean \pm SE. Significant differences between the groups were determined using two-way ANOVA test, with statistical significance set at $p < 0.05$.

RESULTS

Effect of MEPL on cell viability

To determine the effect of MEPL on T24 bladder carcinoma cell viability, an MTT assay was performed 24 h after treatment with the indicated concentrations (50–200 μ g/ml) of MEPL in the absence or presence of TNF- α (20 ng/ml). In the range of 25–150 μ g/ml MEPL showed no cytotoxicity in T24 bladder carcinoma cells. However, the viability of the cells was reduced to

approximately 80 %, when 200 μ g/ml MEPL was used (Fig 1). DMSO (0.1 %) as a solvent (data not shown) in the presence of TNF- α (20 ng/ml) did not produce any cytotoxic effects on the viability of T24 bladder carcinoma cells. Therefore, a 100 μ g/ml concentration of MEPL was used in the remaining experiments.




Fig 1: Effects of MEPL on the viability of T24 bladder carcinoma cells. Cells (1×10^5 cells/ml) were incubated with the indicated concentrations of MEPL (50–200 μ g/ml) 1 h before TNF- α (20 ng/ml) treatment for 24 h. Cell viability was determined by MTT assay; data are mean \pm SE, $n = 3$; * $p < 0.05$

Suppression of MMP-9 gene transcription and its activity by MEPL

In order to assess whether MEPL decreases the expression of MMP-9, zymography, Western blot analysis and RT-PCR were performed. Zymography data showed that MEPL suppresses the expression of MMP-9 activity compared to the TNF- α -stimulated group (Fig 2A). Although MMP-9 protein expression increased significantly in the presence of only TNF- α , Western blot analysis showed decreased MMP-9 protein expression following MEPL pretreatment (Fig 2B).

To assess whether MMP-9 gene expression

is regulated by MEPL, RT-PCR analysis was conducted; the results showed that 6 h of TNF- α treatment significantly increased MMP-9 expression at the transcriptional level (Fig 2C). Furthermore, MMP-9 gene expression was downregulated after pretreatment for 1 h with 100 μ g/ml MEPL, which was similar to the results obtained by zymography and Western blot analysis. Taken together, these data indicate that MEPL suppresses upregulation of TNF- α -induced MMP-9 expression at the transcriptional level.




Fig 2: Effects of MEPL on TNF- α -induced MMP-9 protein and mRNA expression in T24 bladder carcinoma cells. (A) Cells were treated with 100 μ g/ml MEPL 1 h before TNF- α (20 ng/ml) treatment for 24 h. Conditional medium was collected after 24 h, followed by gelatin zymography. (B) Equal amounts of cell lysates were resolved on SDS-polyacrylamide gels, transferred to nitrocellulose membranes, and probed with antibodies against MMP-9. (C) In a parallel experiment, cells (2×10^5 cells/ml) were incubated with the indicated concentrations of MEPL 1 h before TNF- α (20 ng/ml) treatment for 6 h. Total RNA was isolated and RT-PCR was performed using MMP-9 specific primers. **Note:** GAPDH and β -actin were used as internal controls for RT-PCR and Western blot analyses, respectively. The experiments were repeated three times, and similar results were obtained.

Inhibitory effect of MEPL on TNF- α -induced NF- κ B activity

NF- κ B signal pathway is known to play a critical role in modulating MMP-9 expression [12]. The EMSA data showed that TNF- α caused a significant increase in the amount of binding complexes between NF- κ B and specific-binding DNA. Pretreatment with MEPL, however, significantly reduced TNF- α -induced NF- κ B activity in the T24 bladder carcinoma cells (Fig 3A). In a parallel experiment, TNF- α significantly induced phosphorylation of I κ B α and decreased p65 and p50 expression in the cytosol (Fig 3B). However, MEPL sustained p65 and p50




Fig 3: Effect of MEPL on NF- κ B DNA binding activity in T24 bladder carcinoma cells. Cells were preincubated with MEPL (100 μ g/ml) 1 h before stimulation with TNF- α (20 ng/ml) for 30 min. (A) The nuclear extracts were assayed for NF- κ B activity by EMSA and (B) the expression or phosphorylation levels of p65, p50 and p-I κ B α was analyzed using Western blot analysis. **Note:** β -Actin was used as an internal control for western blot analyses. The experiment was repeated three times and similar results were obtained. N.S. = non-specific.

expression and blocked phosphorylation of I κ B α in response to TNF- α . These data indicate that MEPL reduces NF- κ B activity in TNF- α -stimulated T24 bladder carcinoma cells by suppressing I κ B α phosphorylation.

Reduction of invasion by MEPL in T24 bladder carcinoma cells

MMP-9 is thought to be critically involved in the process of tumor invasion [8]. The results of the matrigel assay showed that invasion of T24 bladder carcinoma cells sharply increased, when cells were treated with 20 ng/ml TNF- α (Fig 4A). However, treatment with 100 μ g/ml MEPL 1 h prior to TNF- α treatment significantly reduced the invasion of T24 bladder carcinoma cells to approximately 50 %, as compared to that in the TNF- α -treated group (Fig. 4B). These results indicate that treatment with MEPL inhibits TNF- α -induced invasion of T24 bladder carcinoma cells.




Fig 4: Inhibitory effect of MEPL invasion of T24 bladder carcinoma cells. (A) The upper compartments of Transwells were coated with matrigel for the invasion assay; (B) data are mean \pm SE, n = 3; the number of cells per field were estimated.

DISCUSSION

We previously reported that MEPL has anti-inflammatory effect on LPS-stimulated BV2 microglia cells (unpublished data). However, the mechanism by which MEPL inhibits invasion of tumor cells has not been elucidated completely. Therefore, in this study, we evaluated the effects of MEPL on the invasion of T24 bladder carcinoma cells. Our study is the first to provide evidence that MEPL inhibits TNF- α -induced MMP-9 expression in T24 bladder carcinoma cells by blocking NF- κ B activation.

In the process of cancer invasion, cells can escape from the primary tumor, penetrate into lymphatic and blood vessels, circulate through the bloodstream, and grow into distant foci in normal tissues elsewhere in the body [13]. Cancer cells are involved in numerous interactions with the extracellular matrix and its proteins such as growth factors and cytokines. MMPs are thought to be critical molecules that assist cancer cells during invasion. MMP-9, in particular, is regarded as the main molecule involved in the malignant progression to tumor invasion [14]. MMP-9 activation is specifically associated with tumor progression and invasion in bladder tumors [15]. In particular, recent studies have shown that many extracts isolated from plants attenuate cancer metastasis and angiogenesis via down-regulating the expression of MMP-9 [16-18]. Therefore, inhibition of MMP-9 expression is a strategic target for the development of a therapeutic experimental model of tumor invasion and metastasis. As shown by the matrigel assay, MEPL significantly suppressed cell invasion. According to the zymography data, MEPL treatment resulted in a decrease of MMP-9 activity in the culture medium. These data support the assertion that MEPL is a potential natural resource that can inhibit cancer cell invasion via suppression of MMP-9 expression

Expression of the transcription factor NF- κ B

is induced by various distinct stimuli. Activated NF- κ B, in turn, modulates transcription of many target genes including MMP-9 [16]. NF- κ B is a ubiquitous transcription factor that responds rapidly in mammalian cells and is strongly activated by the cytokines interleukin-1 and TNF- α under various pathological conditions [17,18]. In non-stimulated cells, NF- κ B is present in the cytosol where it is complexed with its inhibitor I κ B. Activation of NF- κ B depends on the phosphorylation of I κ B induced by a signal of a specific I κ B kinase, which initiates conjugation of the inhibitors with ubiquitin and subsequent phosphorylation/degradation [19]. In this study, we showed that MEPL inhibits TNF- α -induced DNA-binding.

CONCLUSION

The results obtained in this study support the fact that MEPL is a tactic regulator that inhibits MMP-9 expression via suppression of NF- κ B activity. We confirmed that MEPL downregulates the expression of MMP-9 activity by suppressing TNF- α -induced activation of NF- κ B. Therefore, MEPL may be an effective therapeutic source for regulating tumor invasion.

ACKNOWLEDGEMENT

This research was supported by the project funded by the Ministry of Land, Transport and Maritime Affairs (grant no. 20088137), Republic of Korea.

COMPETING INTERESTS

The authors report no conflict of interest and they alone are responsible for the contents of this work.

REFERENCES

1. Kanayama H. Matrix metalloproteinases and bladder cancer. *J Med Invest* 2001; 48: 31–43.
2. Landis SH, Murray T, Bolden S, Wingo PA. Cancer statistics. *CA Cancer J Clin* 1999; 49: 8–31.
3. Greenlee RT, Murray T, Bolden S, Wingo PA. Cancer statistics. *CA Cancer J Clin* 2000; 50: 7–33.
4. Chabottaux V, Noel A. Breast cancer progression: insights into multifaceted matrix metalloproteinases. *Clin Exp Metastasis* 2007; 24: 647–656.
5. Egeblad M, Werb Z. New functions for the matrix metalloproteinases in cancer progression. *Nat Rev Cancer* 2002; 2: 161–174.
6. Sier CF, Kubben FJ, Ganesh S, Heerding MM, Griffioen G, Hanemaaijer R, van Krieken JH, Lamers CB, Verspaget HW. Tissue levels of matrix metalloproteinases MMP-2 and MMP-9 are related to the overall survival of patients with gastric carcinoma. *Br J Cancer* 1996; 74: 413–417.
7. Kim S, Choi JH, Kim JB, Nam SJ, Yang JH, Kim JH, Lee JE. Berberine suppresses TNF- α -induced MMP-9 and cell invasion through inhibition of AP-1 activity in MDA-MB-231 human breast cancer cells. *Molecules* 2008; 13: 2975–2985.
8. Rose P, Huang Q, Ong CN, Whiteman M. Broccoli and watercress suppress matrix metalloproteinase-9 activity and invasiveness of human MDA-MB-231 breast cancer cells. *Toxicol Appl Pharmacol* 2005; 209: 105–113.
9. Simon C, Goepfert H, Boyd D. Inhibition of the p38 mitogen-activated protein kinase by SB 203580 blocks PMA-induced Mr 92,000 type IV collagenase secretion and in vitro invasion. *Cancer Res* 1998; 58: 1135–1139.
10. Mineur F, Clerck OD, Roux AL, Maggs CA, Verlaque M. *Polyopes lancifolius* (Halymeniales, Rhodophyta), a new component of the Japanese marine flora introduced to Europe. *Phycologia* 2010; 49: 86–96.
11. Kim KY, Nguyen TH, Kurihara H, Kim SM. α -Glucosidase inhibitory activity of bromophenol purified from the red alga *Polyopes lancifolia*. *J Food Sci* 2010; 75: 145–150.
12. Fidler IJ. Critical factors in the biology of human cancer metastasis: twenty-eighth G.H.A. Clowes memorial award lecture. *Cancer Res* 1990; 50: 6130–6138.
13. Taheri F, Bazan HE. Platelet-activating factor overturns the transcriptional repressor disposition of Sp1 in the expression of MMP-9 in human corneal epithelial cells. *Invest Ophthalmol Vis Sci* 2007; 48: 1931–1941.
14. Pryczynicz A, Guzińska-Ustymowicz K, Dymicka-Piekarska V, Czyżewska J, Kemona A. Expression of matrix metalloproteinase 9 in pancreatic ductal carcinoma is associated with tumor metastasis formation. *Folia Histochem Cytobiol* 2007; 45: 37–40.
15. Torii A, Kodera Y, Uesaka K, Hirai T, Yasui K, Morimoto T, Yamamura Y, Kato T, Hayakawa T, Fujimoto N, Kito T. Plasma concentration of matrix metalloproteinase 9 in gastric cancer. *Br J Surg* 1997; 84: 133–136.
16. Hseu YC, Wu CR, Chang HW, Kumar KJ, Lim MK, Chen CS, Cho HJ, Huang CY, Huang CY, Lee HZ, Hsieh WT, Chung JG, Wang HM, Yang HL.

- Inhibitory effects of Physalis angulata on tumor metastasis and angiogenesis. J Ethnopharmacol 2011; 135: 762-771.*
17. Song TY, Lin HC, Yang NC, Hu ML. Antiproliferative and antimetastatic effects of the ethanolic extract of *Phellinus igniarius* (Linneaus: Fries) Quelet. *J Ethnopharmacol 2008; 115: 50-56.*
 18. Hseu YC, Chen SC, Lin WH, Hung DZ, Lin MK, Kuo YH, Wang MT, Cho HJ, Wang L, Yang HL. *Toona sinensis* (leaf extracts) inhibit vascular endothelial growth factor (VEGF)-induced angiogenesis in vascular endothelial cells. *J Ethnopharmacol 2011; 134: 111-121.*
 19. Miller SC, Huang R, Sakamuru S, Shukla SJ, Attene-Ramos MS, Shinn P, Van Leer D, Leister W, Austin CP, Xia M. Identification of known drugs that act as inhibitors of NF- κ B signaling and their mechanism of action. *Biochem Pharmacol 2010; 9: 1272-1280.*
 20. Epinat JC, Gilmore TD. Diverse agents act at multiple levels to inhibit the Rel/NF- κ B signal transduction pathway. *Oncogene 1999; 18: 6896-6909.*
 21. Karin M, Ben-Neriah Y. Phosphorylation meets ubiquitination: the control of NF- κ B activity. *nu Rev Immunol 2000; 18: 621-663.*
 22. Li Q, Lu Q, Bottero V, Estepa G, Morrison L, Mercurio F, Verma IM. Enhanced NF- κ B activation and cellular function in macrophages lacking I κ B kinase 1 (IKK1). *Proc Natl Acad Sci 2005; 102: 12425-12430.*