

## Original Research Article

# Anti-biofilm and antimicrobial activity of *Mentha pulegium* L essential oil against multidrug-resistant *Acinetobacter baumannii*

Uğur Tutar<sup>1\*</sup>, Cem Çelik<sup>2</sup>, İsa Karaman<sup>3</sup>, Mehmet Ataş<sup>4</sup> and Ceylan Hepokur<sup>5</sup>

<sup>1</sup>Department of Nutrition and Dietetics, Faculty of Health Sciences, <sup>2</sup>Department of Medical Microbiology, Faculty of Medicine, Cumhuriyet University, <sup>3</sup>Department of Bioengineering, Faculty of Natural Sciences and Engineering, Gaziosmanpaşa University, Tokat, <sup>4</sup>Department of Pharmaceutical Microbiology, <sup>5</sup>Department of Biochemistry, Faculty of Pharmacy, Cumhuriyet University, Sivas, Turkey

\*For correspondence: **Email:** [ututar5@gmail.com](mailto:ututar5@gmail.com); **Tel:** +90 346 2192054; **Fax:** +90 346 2191261

Received: 1 February 2016

Revised accepted: 20 April 2016

### Abstract

**Purpose:** To investigate the antimicrobial and anti-biofilm activities of essential oil from *Mentha pulegium* L. (EOMP) on multi-drug resistant (MDR) isolates of *A. baumannii*, as well as its phytochemical composition, antioxidant properties and cytotoxic activity.

**Methods:** The phytochemical composition of EOMP was analyzed by gas chromatography, while its antimicrobial activities were determined by disc diffusion and broth micro-dilution methods. Minimal biofilm inhibition concentration (MBIC) and minimal biofilm eradication concentration (MBEC) tests were used for assessment of its anti-biofilm properties. Viability in the biofilm was studied using 2,3-bis (2-methoxy-4-nitro-5-sulfophenyl)-2H-tetrazolium-5-carboxanilide (XTT) assay, while colorimetric assay was used to assess its cytotoxicity on L929 cells

**Results:** D-isomenthone, pulegone, isopulegone, menthol and piperitenone were the major components of the plant extract. EOMP produced > 22 mm inhibition zone for the isolates, with minimum inhibitory concentration (MIC) and MBIC of 0.6 - 2.5 and 0.6 - 1.25 µL/mL, respectively, while MBEC was ≥ 10 µL/mL. EOMP damaged biofilm structures formed by *A. baumannii* strains at MIC by 26 – 91 %.

**Conclusion:** These results suggest that EOMP contains agents that may be useful in the development of new drugs against *A. baumannii* infections.

**Keywords:** *Mentha pulegium*, Essential oil, Anti-biofilm, Antimicrobial activity, Multi-drug resistance, *A. baumannii*

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

## INTRODUCTION

*Acinetobacter baumannii* is an aerobic, non-fermentative, oxidase-negative and non-motile Gram-negative bacillus. In recent years, the rapidly increasing antimicrobial resistant pattern of this bacterium has become a serious medical problem associated with hospital-acquired infections. *A. baumannii* causes outbreaks of

infection and health care-associated infections, including bacteremia, bloodstream infections, pneumonia, meningitis, urinary tract infection and wound infection [1].

Several factors may be responsible for development of virulence and resistance by *A. baumannii*. One of them is the strong biofilm activity formed by this bacterium. Biofilm

production, which is common in many bacterial species, protects pathogens from antimicrobial agents and the immune responses of their hosts. The frequency and severity *A. baumannii* infections have led to recent and heightened interest in understanding and combating the menace [2].

Sessile microorganisms in the biofilms can be much more resistant to an agent compared to the planktonic forms of the same microorganism. Therefore, it is important to understand the relationship between the biofilm and the resistance encountered in treating bacterial infections [3].

The problem of resistance against the antibiotics, which has been increasing substantially, has led need to search for new and more effective therapeutic agents. Antibiotics were discovered from the rich resources of natural products. The potential of herbal extracts and some of their constituents for overcoming antibiotic-resistant bacteria have been extensively reported [4,5].

*M. pulegium* L. is a flowery plant from the Lamiaceae family which is common worldwide. It is known that this plant (also known as Pennyroyal) is traditionally used for the treatment of many diseases [6]. Although there are publications related to the antimicrobial activity of the *M. pulegium* essential oil (EOMP), not much is known about its anti-biofilm activity. In this study, we aimed at determining the antimicrobial and anti-biofilm effects of the EOMP on multi-drug resistance (MDR) isolates of *A. baumannii*. In addition, the chemical composition of this oil, its antioxidant properties and cytotoxicity were determined.

## EXPERIMENTAL

### Plant material and extraction of EO

The aerial parts of *M. pulegium* were collected in August of 2014 in Manisa flora located in the Aegean Region of Turkey. The samples were collected from Gordes county and dried air-dried at room temperature. The plant was identified and authenticated by a qualified taxonomist in 19 Mayıs University, in Samsun City, in Turkey. At the end of the drying process, the leaves and flowers were ground before the essential oil was extracted by hydro-distillation method using the Clevenger apparatus.

### Analysis and identification of essential oil

The chemical composition of the essential oil was determined according to the method of Aksit

et al [7]. The GC apparatus used was a Perkin-Elmer Clarus 500 Series GC system equipped with a flame ionization detector (FID) and BPX-5 apolar capillary column (30 m x 0.25 mm, 0.25 m i.d.) connected to a mass spectrometer.

### Cultivation of L929 mouse fibroblast cells

Dulbecco's Modified Eagle Medium (DMEM) (Sigma, St. Louis, MO, USA) was used. L929 (ATCC cell line, NCTC clone 929), a mouse connective tissue fibroblast cell line, was cultured in DMEM containing 10 % fetal calf serum (Sigma, St. Louis, MO, USA) and 2 mM/mL L-glutamine. The medium with the cell culture had no antibiotic added. Cultivation of the cultures was performed in an incubator with 5 % CO<sub>2</sub> and at 37 °C until confluence was observed in the cell monolayer, which occurred after about 7 days. At all times, assays were carried out at the exponential growth phase of the cells.

### Cell proliferation assay

This was carried out according to the procedure outlined by Polat et al [8]. In this assay, cell viability is assessed through the cleavage of tetrazolium salts, which are added to the culture medium, and for this, the water soluble tetrazolium (WST-1) labeling reagent (Roche, Mannheim, Germany) was used. Cellular enzymes cleave the tetrazolium salt to formazan, the concentration of which is directly proportional to the number of metabolically active cells in the culture.

Microtiter plates (96-well) were used to seed the cells to a concentration of  $1 \times 10^5$  cells/mL in a 100  $\mu$ L final volume for each well. After 24 h, the cells were treated with different concentrations of EOMP (6.25, 12.5, 25, 50, and 100 % v/v). The cells were then incubated for 24 h at 37 °C in a humidified atmosphere containing 5 % CO<sub>2</sub>. Thereafter, 10  $\mu$ L of culture medium (DMEM) in each well was added to 10  $\mu$ L of the WST-1 labeling reagent. Absorbance was read at 450 nm against control (untreated cells) in a microtiter plate reader (Thermo Scientific Microplate Photometer, Multiskan FC, USA). As a background control, one well contained culture medium and WST-1 labeling reagent (10  $\mu$ L of WST-1 labeling reagent and 10  $\mu$ L of culture medium). The absorbance of this negative control was taken 2 h into the tetrazolium reaction. Cell viability was as in Eq 1.

$$\text{Cell viability (\%)} = (A_s/A_c)100 \dots\dots\dots(1)$$

where  $A_s$  and  $A_c$  are the absorbance of sample and negative control respectively.

### **DPPH (2,2-diphenyl-1-picrylhydrazyl) photometric assay**

The method is based on the reduction of alcoholic solutions of the stable radical, DPPH by electron-donating antioxidants, thereby decolorizing it. The amount of DPPH that is left after a given time is inversely proportional to the radical scavenging activity of the antioxidant [9]. Five mL of 0.004 % solution of DPPH in methanol was added to 50 microliter of EOMP at different concentrations. Absorbance was read against a blank at 517 nm after 30 min of incubation at room temperature [10]. Eq 2 was used to calculate inhibition of DPPH.

$$\text{Inhibition (\%)} = \{(A_0 - A_1)/A_0\}100 \dots\dots\dots(2)$$

where A0 = control absorbance and A1 = sample absorbance. Ascorbic acid, a natural antioxidant was used as positive control. All the tests were conducted in triplicate.

### **Microorganisms**

*A. baumannii* strains used in the study were isolated in the Clinical Microbiology Laboratories of Cumhuriyet University Hospital between 2013 and 2014. The bacteria were identified using the BD Phoenix (Becton Dickinson, Sparks, MD, USA) automated microbiology system and antimicrobial susceptibility tests were conducted. Resistance and susceptibility of the strains against the antimicrobial agents were assessed according to the Clinical Laboratory Standards Institute (CLSI) [11].

### **Evaluation of antimicrobial and anti-biofilm activity**

#### **Disc diffusion assay**

Disc diffusion method was applied, with 100 µL of suspension containing 10<sup>8</sup> CFU/mL of bacteria on Mueller Hinton Agar (MHA). After impregnation of the disc (6 mm in diameter) with 20 µL of EOMP (10 µL/disc) at a concentration of 500 µl/mL, it was placed on the inoculated agar. The solvent used for dissolving EOMP was utilized to prepare the negative control. The inoculated plates were incubated at 37 °C for 24 h. Zone of inhibition was measured to assess the antimicrobial activity against each test organism. All the assays were repeated three times.

#### **Microdilution assay**

The broth micro-dilution method which was employed to assess the antimicrobial activities of

the essential oil was done in line with the protocol of the CLSI [11]. A micro-well dilution method was taken as a basis for determining the MIC of EOMP against the bacterial strains.

Twenty four-hour broth cultures were used to prepare the inoculums of the microorganisms, and suspensions were adjusted to 0.5 McFarland standard turbidity. EOMP was dissolved in Mueller Hinton Broth (MHB) containing 0.5 % (v/v) Tween 20, and serial two-fold dilutions were prepared in 96-well plates with MHB at a concentrations varying between 0.03 and 2 % (v/v). Whereas there was MHB and cells without EOMP in the wells of the positive control, wells of negative control had only MHB. A BioTek microplate reader (Bio Tek Instruments, Inc., Winooski, VT, USA) was used to measure the absorbance of the plates at 550 nm before and after incubation for 24 h at 37 °C. The minimum inhibitory concentration (MIC) was detected as the lowest concentration of EOMP, and at this concentration, the optical density at the 24th h incubation of the inoculum remained the same or decreased when compared to the reading at the beginning.

Ten µL was taken from each well following the incubation, spot-inoculated on MHA and incubated at 37 °C for 24 h in order to determine the minimum bactericidal concentration (MBC) of EOMP. The concentration without any growth of the subculture was taken as the MBC.

### **Minimum biofilm inhibitory concentration (MBIC) assay**

Minimum biofilm inhibitory concentration (MBIC) is described as the lowest concentration at which the antimicrobial agents prevents the biofilm formation [12]. In order to test the anti-biofilm activity of EOMP against *A. baumannii*, the microtiter plate assay was used.

The MBIC test was carried out using the method of Adukwu *et al* [13]. An aliquot (100 µL) from an overnight culture (diluted with MHB containing 1 % (w/v) glucose to 10<sup>8</sup> CFU/ mL) was dispensed into each test well of a 96-well plate. Then 100 µL of different concentrations of EOMP (0.03 – 2 %) were dispensed into different wells. The negative control contained only MHB whereas the positive control contained cell cultures alone without EOMP. The supernatant of the wells were decanted and each well gently rinsed three times with 300 µL of sterile distilled water and discarded after incubation at 37 °C for 24 h. The plates were dried in air for 30 min and stained with 0.1 % (w/v) crystal violet at room temperature for 30 min, washed 3 times with

distilled water and dried. Thereafter the crystal violet was solubilized in 95 % ethanol and absorbance was read in a microplate reader (Bio Tek Instruments, Inc., Winooski, VT, USA) at 550 nm. MBIC was determined as the EOMP concentration at which the absorbance is equal to or less than that of the negative control. This test was performed in triplicate and the mean ( $n = 3$ ) was taken.

### Minimum biofilm eradication concentration (MBEC) assay

Minimum biofilm eradication concentration (MBEC) is the lowest concentration which can damage the structure of emerging biofilm [12]. MBEC assay was carried out according to the method of Kuzma *et al* [14]. In essence, 200  $\mu\text{L}$  ( $10^8$  CFU/mL) of each strain was inoculated into each well of flat bottom 96-well microtiter plate. The wells were incubated at 37 °C for 48 h. After biofilm formation, the wells were washed four times with sterile distilled water in order to remove non-adherent cells. Serial dilutions of EOMP (0.03 – 2 %) were added to different wells (each EOMP concentration at a volume of 200  $\mu\text{L}$ ). Incubation of the plates was at 37 °C for 24 h and thereafter the wells were washed with distilled water and stained with crystal violet. Biofilm without EOMP served as positive control. The concentration at which established biofilms were removed from the bottom of the treated wells was taken as the MBEC.

### Evaluation of biofilm eradication

Each *A. baumannii* strain (200  $\mu\text{L}$  of  $10^8$  CFU/mL) was inoculated into each well of flat bottom 96-well microtiter plate and incubated for 48 h at 37 °C. After biofilm formation, EOMP was added to different wells at concentrations corresponding to MIC, 1/2MIC, 1/4 MIC and 1/8 MIC values. The wells were incubated for 24 h, washed twice with phosphate buffer saline and read at 550 nm. Biofilm eradication was calculated as in Eq 3.

Biofilm eradication (%) =  $\{(Ac - As)/Ac\}100 \dots (2)$  [15]

where Ac and As are the absorbance of sample and control respectively.

### Biofilm metabolism assay

Evaluation of the effect of EOMP on the metabolic activity of the biofilms formed by the bacterial strains was carried out through a modification of the tetrazolium salt (XTT) reduction test, according to the method of Chaieb *et al* [15]. EOMP at concentrations of 1.25 and

2.5  $\mu\text{L}/\text{mL}$  was applied to the 48 h biofilm structures formed in 96-well plates, and after 24 h incubation, XTT reagent was added to the test and control wells. At the end of 5 h incubation period, cell viability was determined by reading the wells at 450 nm in a microplate reader.

### Statistical analysis

The results are presented as mean  $\pm$  standard deviation ( $n = 3$ ) and assessed by one-way analysis of variance (ANOVA), while Student's t-test was applied for comparison between two groups.  $P < 0.05$  were considered statistically significant.

## RESULTS

### GC and GC-MS data

Table 1 shows results from GC and GC-MS analysis of EOMP.

**Table 1:** Main components of essential oil of aerial parts of *Mentha pulegium*

Rt <sup>a</sup>	Compound	(%) <sup>b</sup>
4.407	p-Xylene	0.15
5.689	$\alpha$ -Pinene	0.06
6.026	Cyclohexanone, 3-methyl-	0.10
6.661	$\beta$ -pinene	0.06
7.096	3-Octanol	0.32
10.489	3-Octanol, acetate	0.32
11.163	$\alpha$ -Terpinolen	0.10
11.299	Isopulegone	9.20
11.587	D-isomenthone	33.19
11.899	Menthol	2.98
13.729	Pulegone	51.32
14.042	Piperitone	0.83
16.348	Piperitenone	0.86


Rt<sup>a</sup> = retention time; (%)<sup>b</sup> = relative percentage obtained from peak area

### Cytotoxic potential of *M. pulegium* EO


Results of cytotoxicity potential of EOMP are shown in Figure 1. The toxicity potential of the 50 % and 100 % EOMP were statistically different from that of the control (Figure 1,  $p < 0.05$ ). However there were no significant differences in toxicity potential between the control and EOMP at concentrations of 6.25, 12.5 and 25 %.

### Antioxidant capacity of EOMP

Figure 2 shows the antioxidant capacity of EOMP with respect to DPPH scavenging. The EOMP scavenged DPPH with an IC<sub>50</sub> of  $8.82 \pm 0.67$   $\mu\text{g}/\text{mL}$ , compared to that of ascorbic acid with IC<sub>50</sub> of  $6.97 \pm 1.05$   $\mu\text{g}/\text{mL}$ .



**Figure 1:** *In vitro* cytotoxic effect of EOMP on L929 fibroblast cells. \* $p < 0.05$  compared to control



**Figure 2:** DPPH radical scavenging activity of EOMP.  $IC_{50}$  is the concentration of EOMP that scavenged 50 % of DPPH radical. All data are presented as the mean  $\pm$  SD for three replicates

## Antimicrobial and antibiofilm activities

The inhibition zones due to antimicrobial effect of EOMP on the 22 strains of *A. pulegium* tested are shown in Table 2. EOMP formed more than 22 mm inhibition zones in all the strains. The MIC values of EOMP were 2.5  $\mu\text{L}/\text{mL}$  in 15 strains, 1.25  $\mu\text{L}/\text{mL}$  in 2 strains, and 0.6  $\mu\text{L}/\text{mL}$  in 5 strains. Values of MBC were 5  $\mu\text{L}/\text{mL}$  in 18 strains and 2.5  $\mu\text{L}/\text{mL}$  in 4 strains. MBIC value was 0.6  $\mu\text{L}/\text{mL}$  in 2 strains, and 1.25  $\mu\text{L}/\text{mL}$  in the other strains. MBEC value was  $\geq 10$   $\mu\text{L}/\text{mL}$  in all the strains.

## Reduction in biofilm formation

MIC, 1/2MIC, 1/4 MIC and 1/8 MIC values of the *M. pulegium* EO were applied onto the bacterial biofilm formations that emerged after the 48h incubation. It was observed that the effect of the EO on the biofilm structure decreased as the concentration reduced. It was seen that the *M. pulegium* EO damaged the biofilm structure at the MIC value by 26-91 % (Table 3).

## Inhibition of metabolic activity

EOMP decreased the metabolic activity of the 48 h biofilm structures formed by MDR *A. baumannii* strains in a concentration-dependent manner (Figure 3).

**Table 2:** Antimicrobial and anti-biofilm activities of EOMP against MDR *Acinetobacter baumannii*


Strain	Zone of inhibition <sup>a</sup>	MIC ( $\mu\text{L}/\text{mL}$ )	MBC ( $\mu\text{L}/\text{mL}$ )	MBIC ( $\mu\text{L}/\text{mL}$ )	MBEC ( $\mu\text{L}/\text{mL}$ )
1	24.6 $\pm$ 2.0	2.5	5	1.25	$\geq 10$
2	26.3 $\pm$ 3.0	2.5	5	1.25	$\geq 10$
3	27.6 $\pm$ 1.1	2.5	5	1.25	$\geq 10$
4	23.3 $\pm$ 1.1	2.5	5	1.25	$\geq 10$
5	22.0 $\pm$ 1.5	2.5	5	1.25	$\geq 10$
6	22.0 $\pm$ 1.5	2.5	5	1.25	$\geq 10$
7	22.6 $\pm$ 0.8	2.5	5	1.25	$\geq 10$
8	22.6 $\pm$ 0.8	2.5	5	1.25	$\geq 10$
9	30.6 $\pm$ 3.7	0.6	2.5	1.25	$\geq 10$
10	31.1 $\pm$ 3.0	0.6	2.5	0.6	$\geq 10$
11	22.6 $\pm$ 0.8	2.5	5	1.25	$\geq 10$
12	24.3 $\pm$ 1.1	2.5	5	1.25	$\geq 10$
13	32.6 $\pm$ 0.5	0.6	2.5	0.6	$\geq 10$
14	23.0 $\pm$ 3.0	2.5	5	1.25	$\geq 10$
15	23.6 $\pm$ 1.5	2.5	5	1.25	$\geq 10$
16	27 $\pm$ 1.7	1.25	5	1.25	$\geq 10$
17	31.6 $\pm$ 3.5	0.6	2.5	1.25	$\geq 10$
18	23.3 $\pm$ 1.1	0.6	5	1.25	$\geq 10$
19	25.3 $\pm$ 2.5	2.5	5	1.25	$\geq 10$
20	25.6 $\pm$ 2.0	2.5	5	1.25	$\geq 10$
21	26.3 $\pm$ 1.5	2.5	5	1.25	$\geq 10$
22	27.6 $\pm$ 1.5	1.25	5	1.25	$\geq 10$

MIC = minimum inhibitory concentration, MBC = minimum bactericidal concentration, MBIC = minimum biofilm inhibitory concentration, MBEC = minimum biofilm eradication concentration; <sup>a</sup>values are reported as mean  $\pm$  SD (mm, n = 3)

**Table 3:** Reduction in biofilm formation

Strain	MIC	1/2 MIC	1/4 MIC	1/8 MIC
1	85.6±1.5	56.3±3.05	25.0±1.7	8.3±1.5
2	85.0±4.3	84.3±4.7	23.3±2.5	2.3±1.1
3	86.0±7.0	89.3±1.1	16.3±3.5	5.3±2.5
4	86.6±2.0	33.3±2.5	34.0±2.0	12.6±0.5
5	80.3±0.5	75.6±3.0	25.0±3.6	13.6±2.0
6	87.6±1.5	82.6±4.5	18.3±4.0	2.3±1.5
7	89.0±1.0	84.3±5.6	27.6±3.5	4.6±3.7
8	78.0±1.7	77.0±4.0	13.6±1.1	2.6±0.5
9	26.3±2.0	5.0±1.0	NT	NT
10	41.0±2.6	2.33±0.5	NT	NT
11	91.0±2.6	90.3±2.5	28.6±0.5	24.0±2.0
12	87.0±2.0	35.0±2.6	22.6±1.5	9.6±3.5
13	36.0±2.6	8.3±1.5	NT	NT
14	84.6±3.2	78.0±8.1	25.0±5.2	25.3±2.5
15	82.0±4.0	73.0±6.2	11.3±1.5	10.3±3.2
16	56.3±6.1	26.3±7.5	24.0±3.6	NT
17	57.3±2.8	46.6±4.04	NT	NT
18	34.3±4.1	25.6±3.05	NT	NT
19	83.6±5.1	76.6±6.5	32.6±1.5	3.0±1.7
20	84.3±6.02	79.6±4.7	47.6±3.5	38.0±3.0
21	86.3 ±0.5	73.0±2.0	46.6±5.5	26.6±3.2
22	83.3±3.5	69.0±2.6	33.3±2.8	NT

Data are reported as mean ± SD (n = 3); NT =not tested



**Figure 3:** Changes in metabolic activity following 48 h exposure of biofilms of MDR *A. baumannii* strains to EOMP, as determined by XTT assay (control = biofilms not exposed to EOMP). All data are presented as mean ± SD for three replicates

## DISCUSSION

Natural products are sources of chemical compounds that have been used as alternative medicines to conventional therapy. Currently, researchers are focusing on the pharmacological and therapeutic activities of plant products.

This study demonstrates that the essential oil of *Mentha pulegium* exhibits anti-biofilm and antimicrobial activities multi-drug resistant strains of *Acinetobacter baumannii*. In addition, the

study demonstrated the antioxidant properties, cytotoxic activity and chemical composition of *M. pulegium* EO.

*M. pulegium* essential oil has strong antimicrobial activity on MDR *A. baumannii* isolates. It was observed that the MIC which has effect on planktonic forms are not adequately effective on the biofilm structure. In the XTT test, MIC was not completely effective on the sessile forms in the biofilm, so that viability continued. Therefore, we concluded that it will be appropriate to

investigate the anti-biofilm activity of the essential oil, and regulate the treatments accordingly. All the *A. baumannii* strains used in this study were found resistant to the tested antibiotics (amikacin, cefepime, ceftazidime, gentamycin, imipenem, meropenem). The cytotoxic potential of 50 % EOMP was significantly different from that of control. The MIC, MBIC and MBEC values of EOMP ( $\geq 0.6$ - $2.5$   $\mu\text{l/mL}$ ,  $0.6$ - $2.5$   $\mu\text{l/mL}$  and  $10$   $\mu\text{l/mL}$ , respectively) were quite below the cytotoxic value, blocked microbial biofilm formation even at low concentrations, and eradicated emerging biofilm structures in a concentration-dependent manner even at concentrations below MIC. These results indicate that EOMP has effective antimicrobial and anti-biofilm activities against *A. baumannii*.

DPPH radical scavenging activity assay was used to evaluate the possible antioxidant properties of the *M. pulegium* essential oil. The results obtained confirm previous reports on the importance of essential oils as natural antioxidants [23-25]. A previous study reported that extracts from some *Mentha* species (*M. longifolia* (L.) Huds., *M. piperita* L., *M. pulegium* L., *M. rotundifolia* (L.) Huds., and *M. spicata* L.) have antioxidant and free radical -scavenging properties [26]. The antioxidant properties of *Mentha* extracts may be due to their phytochemical compositions. Previous studies reported that *M. pulegium* has strong antioxidant components [27]. According to result of the present study, *M. pulegium* essential oil can be considered as a strong antioxidant and a strong antimicrobial agent.

The major components of EOMP were pulegone (51.32 %), D-isomenthone (33.19 %), isopulegone (9.20 %), menthol (2.98 %) and piperitenone (0.86 %). This is in agreement with components reported from other studies on *M. pulegium* [28,29]. In 2005, Agnihotri *et al* reported pulegone (65.9-83.1 %) and menthone (8.3-8.7 %) as the major components of the oil of *M. pulegium* collected from different locations in India. Pulegone, piperitenone and isomenthone were reported as the main compounds of the oil of *M. pulegium* from Bulgaria [30,31]. It was also stated in these studies that the essential oil had antimicrobial activity against various microorganisms. The major components of the oil were similar, but may vary according to geographical regions where the plant was harvested, or due to seasonal conditions [23,28,29,32]. Some previous studies reported that pulegone, isomenthone, piperitenone exhibited high antibacterial and antifungal activities [21,33,34]. Thus the antimicrobial and

anti-biofilm properties of EOMP could be attributed to its chemical components.

## CONCLUSION

The findings of this study show that *M. pulegium* essential oil has strong antimicrobial and anti-biofilm activity against MDR *A. baumannii* isolates. Thus, it is a potential agent for the treatment of MDR *A. baumannii* infections. In addition, the results provide evidence that the essential oil of *M. pulegium* may be a useful source of antioxidants.

## ACKNOWLEDGEMENT

The authors would like to thank Cumhuriyet University Faculty of Medicine microbiology laboratory staff for assistance in this research.

## CONFLICT OF INTEREST

No conflict of interest associated with this work.

## CONTRIBUTION OF AUTHORS

We declare that this work was done by the authors named in this article and all liabilities pertaining to claims relating to the content of this article will be borne by the authors.

## REFERENCES

- Howard A, O'Donoghue M, Feeney A, Sleator RD. *Acinetobacter baumannii*: an emerging opportunistic pathogen. *Virulence* 2012; 3(3): 243-250.
- Aşık G. Current approaches to explain the virulence of *Acinetobacter baumannii*. *Mikrobiyol Bul* 2011; 45(2): 371-380.
- Penesyan A, Gillings M, Paulsen IT. Antibiotic discovery: Combatting bacterial resistance in cells and in biofilm communities. *Molecules* 2015; 20(4): 5286-5298.
- Yap PS, Yiap BC, Ping HC, Lim SH. Essential oils a new horizon in combating bacterial antibiotic resistance. *Open Microbiol J* 2014; 8(Feb): 6-14.
- Koehn FE. High impact technologies for natural products screening. *Prog Drug Res* 2008; 65: 175, 177-210.
- Ak I. New terpenoids from *Mentha pulegium* and their antimicrobial activity. *Nat Prod Res* 2013; 27(8): 691-696.
- Aksit H, Demirtas I, Telci I, Tarimcilar I. Chemical diversity in essential oil composition of *Mentha longifolia* (L.) Hudson subsp. *typhoides* (Briq.) Harley var. *typhoides* from Turkey. *J Essent Oil Res* 2013; 25(5): 430-437.
- Polat ZA, Walochnik J, Obwaller A, Vural A, Dursun A, Arici MK. Miltefosine and polyhexamethylene biguanide: a new drug combination for the treatment of

- Acanthamoeba keratitis*. *Clin Experiment Ophthalmol* 2014; 42(2): 151-158.
9. Burits M, Bucar F. Antioxidant activity of *Nigella sativa* essential oil. *Phytother Res* 2000; 14(5): 323-328.
  10. Mensor LL, Menezes FS, Leitão GG, Reis AS, dos Santos TC, Coube CS, Leitão SG. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. *Phytother Res* 2001; 15(2): 127-130.
  11. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fourth Informational Supplement. CLSI document M100-S24. Wayne, PA (2014).
  12. Ceri H, Olson ME, Morck DV, Storey DG. Minimal biofilm eradication (MBEC) assay: susceptibility testing for biofilms. In: Pace JL, Rupp ME, Finch RG, editors. *Biofilms, infection, and antimicrobial therapy*. CRC Pres, Boca Raton, 2006, pp. 257-269.
  13. Adukwu EC, Allen SC, Phillips CA. The anti-biofilm activity of lemongrass (*Cymbopogon flexuosus*) and grapefruit (*Citrus paradisi*) essential oils against five strains of *Staphylococcus aureus*. *J Appl Microbiol* 2012; 113(5): 1217-1227.
  14. Kuzma L, Rozalski M, Walencka E, Rozalska B, Wysokinska H. Antimicrobial activity of diterpenoids from hairy roots of *Salvia sclarea* L: salvipisone as a potential anti-biofilm agent active against antibiotic resistant *Staphylococci*. *Phytomedicine* 2007; 14(1): 31-35.
  15. Chaieb K, Kouidhi B, Jrah H, Mahdouani K, Bakhrouf A. Antibacterial activity of Thymoquinone, an active principle of *Nigella sativa* and its potency to prevent bacterial biofilm formation. *BMC Complement Altern Med* 2011; 11(Apr): 29.
  16. Peleg AY, Seifert H, Paterson DL. *Acinetobacter baumannii*: emergence of a successful pathogen. *Clin Microbiol Rev* 2008; 21(3): 538-582.
  17. Manchanda V, Sanchaita S, Singh NP. Multidrug resistant *Acinetobacter*. *J Glob Infect Dis* 2010; 2(3): 291-304.
  18. Prakasam G, Bhashini M; Lakshmi Priya, Ramesh SS. In-vitro antibacterial activity of some essential oils against clinical isolates of *Acinetobacter baumannii*. *Indian J Med Microbiol* 2014; 32(1): 90-91.
  19. Jazani NH, Zartoshti M, Babazadeh H, Ali-daiee N, Zarrin S, Hosseini S. Antibacterial effects of Iranian fennel essential oil on isolates of *Acinetobacter baumannii*. *Pak J Biol Sci* 2009; 12(9): 738-741.
  20. Saghi H, Bahador A, Khaledi A, Kachoei RA, Dastjerdi FA, Esmaeili D. Antibacterial effects of *Origanum vulgare* essence against multidrug-resistant *Acinetobacter baumannii* isolated from selected hospitals of Tehran, Iran. *Avicenna J Clin Microb Infec* 2015; 2(1): e22982.
  21. Mahboubi M, Haghi G. Antimicrobial activity and chemical composition of *Mentha pulegium* essential oil. *J Ethnopharmacol* 2008; 119(2): 325-327.
  22. Ceri H, Olson ME, Stremick C, Read RR, Morck D, Buret A. The calgary biofilm device: new technology for rapid determination of antibiotic susceptibilities of bacterial biofilms. *J Clin Microbiol* 1999; 37(6): 1771-1776.
  23. Teixeira B, Marques A, Ramos C, Batista I, Serrano C, Matos O, Neng NR, Nogueira JMF, Saraiva JA, Nunes ML. European pennyroyal (*Mentha pulegium*) from Portugal: Chemical composition of essential oil and antioxidant and antimicrobial properties of extracts and essential oil. *Ind Crop Prod* 2012; 36(1): 81-87.
  24. Cherrat L, Espina L, Bakkali M, Pagán R, Laglaoui A. Chemical composition, antioxidant and antimicrobial properties of *Mentha pulegium*, *Lavandula stoechas* and *Satureja calamintha* Scheele essential oils and an evaluation of their bactericidal effect in combined processes. *Innov Food Sci Emerg* 2014; 22(April): 221-229.
  25. Hajlaoui H, Trabelsi N, Noumi E, Snoussi M, Fallah H, Ksouri R, Bakhrouf A. Biological activities of the essential oils and methanol extract of two cultivated mint species (*Mentha longifolia* and *Mentha pulegium*) used in the Tunisian folkloric medicine. *World J Microbiol Biotechnol* 2009; 25(12): 2227-2238.
  26. Nickavar B, Alinaghi A, Kamalinejad M. Evaluation of the antioxidant properties of five *Mentha* species. *Iran J Pharm Res* 2008; 7(3): 203-209.
  27. Zaidi F, Voirin B, Jay M, Viricel MR. Free flavonoid aglycones from leaves of *Mentha pulegium* and *Mentha suaveolens* (Labiatae). *Phytochemistry* 1998; 48(6): 991-994.
  28. Ghazghazi H, Chedia A, Weslati M, Trakhna F, Houssine S, Abderrazak M, Brahim H. Chemical composition and in vitro antimicrobial activities of *Mentha pulegium* leaves extracts against foodborne pathogens. *J Food Safety* 2013; 33(3): 239-246.
  29. Kanakis CD, Petrakis EA, Kimbaris AC, Pappas C, Tarantilis PA, Polissiou MG. Classification of Greek *Mentha pulegium*. (Pennyroyal) samples, according to geographical location by Fourier transform infrared spectroscopy. *Phytochem Anal* 2012; 23(1): 34-43.
  30. Agnihotri VK, Agarwal SG, Dhar PL, Thappa RK, Baleshwar Kapahi BK, Saxena RK, Qazi GN. Essential oil composition of *Mentha pulegium* L. growing wild in the North-Western Himalayas India. *Flavour Fragr J* 2005; 20(6): 607-610.
  31. Stoyanova A, Georgiev E, Kula J, Majda T. Chemical composition of the essential oil of *Mentha pulegium* L. From Bulgaria. *J Essent Oil Res* 2005; 17(5): 475-476.
  32. Shahmohamadi R, Sariri R, Rasa M, Aghamali M. Antioxidant activity of Gilan *Mentha pulegium* during growth. *Pak J Biol Sci* 2014; 17(3): 380-387.
  33. Primo V, Rovera M, Zanon S, Oliva M, Demo M, Daghero J, Sabini L. Determination of the antibacterial and antiviral activity of the essential oil from *Mintostachys verticillata* (Griseb.) Epling *Rev Argent Microbiol* 2001; 33(2): 113-117.
  34. Flamini G, Cioni PL, Puleio R, Morelli I, Panizzi L. Antimicrobial activity of the essential oil of *Calamintha*



*nepeta and its constituent pulegone against bacteria and fungi. Phytother Res 1999; 13(4): 349-351.*