

## Original Research Article

# Luteolin as a potent anti-leishmanial agent against intracellular *Leishmania tropica* parasite

Kashif Iqbal<sup>1,3\*</sup>, Qaiser Jamal<sup>4</sup>, Javeid Iqbal<sup>1</sup>, Maria Sadaf Afreen<sup>5</sup>, Muhammad Zeeshan Ahmed Sandhu<sup>2</sup>, Eshwa Dar<sup>2</sup>, Umar Farooq<sup>2</sup>, Mohammad Fahd Mushtaq<sup>2</sup>, Numera Arshad<sup>2</sup> and Muhammad Mohsin Iqbal<sup>2</sup>

<sup>1</sup>Department of Pharmacology, Faculty of Pharmacy and Health Sciences, University of Balochistan, Quetta, <sup>2</sup>Islam College of Pharmacy, Sialkot, Punjab, <sup>3</sup>School of Pharmacy, University of Lahore-Islamabad Campus, Islamabad, Pakistan. <sup>4</sup>Department of Zoology, University of Peshawar, Peshawar, <sup>5</sup>Gomal University, Dera Ismail Khan, KP, Pakistan

\*For correspondence: **Email:** [kashifiqbal321@gmail.com](mailto:kashifiqbal321@gmail.com); **Tel:** +92 3356951284

Received: 1 November 2016

Revised accepted: 16 January 2017

### Abstract

**Purpose:** To examine the anti-leishmanial and cytotoxic effects of five naturally occurring phenolic compounds: luteolin (1), lalioside (2), luteolin-4'-O- $\beta$ -D-glucopyranoside (3), apigenin 4-O- $\beta$ -D-glucopyranoside (4) and apigenin (5) on *Leishmania tropica* KWH23 amastigotes.

**Methods:** The compounds were isolated from the leaves of *Lawsonia Inermis* via hyphenated high performance liquid chromatography-high resolution mass spectrometry coupled with solid phase extraction-tube transfer nuclear magnetic resonance technique. The isolated compounds were given intraperitoneally to *L. tropica* KWH23 amastigotes-infected albino mice at a dose of  $\geq 3$  mg/kg for 5 days. Amphotericin-B was used as standard (reference) drug. Lymphocytes were used to analyze their cytotoxicity.

**Results:** For compound 1, mean lesion size decreased from  $0.82 \pm 0.12$  to  $0.10 \pm 0.01$  after 120 days, with 97 % cure of intracellular *L. tropica* amastigotes at a dose of 15 mg/kg, compared to amphotericin B which produced 95 % cure at a dose of 30 mg/kg. Half-maximal concentration ( $IC_{50}$ ) for compound 1 was 4.15  $\mu$ g/ml against lymphocytes.

**Conclusion:** The results indicate that luteolin is a potent inhibitor of *L. tropica* amastigotes, with a higher cytotoxic activity against lymphocytes, compared with luteolin-4'-O- $\beta$ -D-glucopyranoside.

**Keywords:** *Leishmania tropica*, Luteolin, Lalioside, Luteolin-4'-O- $\beta$ -D-glucopyranoside, Apigenin 4-O- $\beta$ -D-glucopyranoside, Apigenin

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

## INTRODUCTION

Leishmaniasis, caused by parasites belonging to the genus *Leishmania* (Family Trypanosomatidae), is a major public health problem in tropical and sub-tropical regions. The parasite is transmitted by the sand fly (vector), with dogs, sheep, rats, horses, and cats being common animal hosts. The World Health Organization, WHO has reported that people from 98 countries

in 5 continents, are at high risk of leishmaniasis, and it is estimated that approximately 12 million people are currently infected [1-3]. Cutaneous leishmaniasis is caused by different *Leishmania* species, e.g. *Leishmania tropica*, *Leishmania major*, *Leishmania amazonensis*, and *Leishmania brazillensis*. In Pakistan, *L. tropica* and *L. major* are the main causes of cutaneous leishmaniasis [4,5]. First line therapy for cutaneous leishmaniasis in Europe, Asia and

Africa is pentavalent antimonials, *i.e.*, sodium stibogluconate and meglumine antimoniate [2]. However, antimonials have severe side effects like myalgia, pancreatitis, cardiac arrhythmia, hepatitis, and drug accumulation in liver and spleen. Thus, there is an urgent need for new chemical entities for non-toxic and effective treatment of leishmaniasis [2,4].

Phenolic compounds comprise different groups including coumarins, flavonoids, tannins, naphthalenes, naphthoquinones, xanthenes, lignans and alkylphenones [6]. Flavonoids and alkylphenones possess a variety of biological and pharmacological activities, including antioxidant [7], antibacterial, antiulcer [8], antifungal [9], antiviral [10], HIV - inhibiting [11], and anti-herpes simplex type 1 (HSV-1) [12]. They also inhibit electron transfer in mitochondrial inner membrane [13]. Flavonoids and alkylphenones are chemically well investigated, and a large number of member compounds have been reported. These include flavonoids and luteolin [14]; luteolin-4'-O- $\beta$ -D-glucopyranoside [15]; apigenin 4-O- $\beta$ -D-glucopyranoside [16]; apigenin [15]; xanthoangelol [8]; macaranggin [7]; artelastin [10]; 5,7-Dihydroxy-6,8-diphenyl flavonoids [11]; leachianone G [12] and alkylphenones such as Laliolide [17], 2,4,6-trihydroxyacetophenone-2-

O- $\beta$ -D-glucopyranoside [18] and lawsoniaside [19].

However, the current work deals with anti-leishmanial and cytotoxic effects of naturally occurring plant phenolic compounds having four flavonoids and one alkylphenone.


## EXPERIMENTAL

### Chemicals

Fetal bovine serum, dimethylsulphoxide, Luteolin, Laliolide, luteolin-4'-O- $\beta$ -D-glucopyranoside, apigenin 4-O- $\beta$ -D-glucopyranoside, apigenin and RPMI-1640 medium were purchased from Sigma-Aldrich (St. Louis, MO), whereas penicillin and streptomycin were purchased from MERCK, U.S.A. Amphotericin B was supplied by AmB-Fungizone, Bristol-Myers Squibb, UK. Water used for *in vivo* test solutions was purified by deionization and 0.22  $\mu$ m membrane filtration (Millipore, Billerica, MA).

### *In vivo* test

To study the pathogenesis of *leishmania* strain, 9 groups of male BALB/c mice (aged 6 - 8 weeks, and weighing 20 - 32 g) were used. Drug administration was through cardiac route.



**Figure 1:** Chemical structures of anti-leishmanial phenolic compounds: Luteolin (1), Laliolide (2), Luteolin-4'-O- $\beta$ -D-glucopyranoside (3), Apigenin 4-O- $\beta$ -D-glucopyranoside (4) and Apigenin (5)

Promastigotes of *L. Tropica* KWH23 [21] were cultured in RPMI-1640 medium along with 10 % fetal bovine serum, penicillin (200 U/ml) and streptomycin (0.2 mg/ml). The parasite was cultured at 26 °C for 4 days in BOD incubator (Gallenkamp, Size 1, UK), and then harvested parasite. The harvested promastigotes were taken in a sterile tube and counted in a haemocytometer (REICHERT, N.Y, USA) under upright microscope (CX31, OLYMPUS, Tokyo, Japan). The promastigotes were centrifuged at 4 °C for 10 min at 2000 rpm; the supernatant liquid was discarded while pellet was left in tube. Fresh RPMI-1640 medium with 10 % FBS was added to get the required volume (10ml). The required volume (10 µl) of promastigotes (containing  $1.4 \times 10^6$  promastigotes/ml) was injected into the cardiac cavity (intraperitoneally) of the BALB/c mice. Developed lesions were measured weekly with dial micrometer (Mitutoyo, Japan) during the infection period. Infection was well established and clearly visible lesions were evident to the naked eye after 36 days. Then treatment process was started. Dose of test compounds given to Groups I, II, III, IV and V were 3.0 mg/kg for 5 days (total dose = 15 mg/kg) in DMSO up to final volume of 3 ml. Amphotericin-B was used as standard drug (positive control) at a dose of 15 mg/kg. No drug agent was used in VII group (negative control). The injection dose of 10 µl was given five times with 3-day intervals, and lesion measurements were recorded regularly. Dial micrometer was used to note the difference between size of the lesion in infected and uninfected mice weekly. Before and after treatment, needle aspirations were taken from the lesions [22]. To detect amastigotes under upright microscope, Giemsa stain was used, and the samples were examined under oil immersions. On the 30<sup>th</sup>, 60<sup>th</sup>, 90<sup>th</sup> and 120<sup>th</sup> day of infection, 60 mg of tissue sample was taken from the lesion for biopsy. In the identification of amastigotes, sample was smeared on the slides stained with Giemsa and upright microscope was used for examination.

### Ethics statement

BALB/c mice were supplied by Department of Pharmacology (Animal center), University of Peshawar, KPK, Pakistan and this study was approved by Animal and Ethics Committee, Faculty of Pharmacy and Health Sciences, University of Balochistan (UOB), Quetta (approval ref. no. 093/FOPHS/UOB). The animals were maintained in accordance with UOB, Quetta Policy and international guidelines on the care and use of laboratory animals [20]. Standard diet along with water were given *ad libitum* to the BALB/c mice during experiments.

### Cytotoxicity test

Fresh blood (10 ml) from a healthy volunteer was taken in BD vacutainer K2E (EDTA) to get mammalian cells (lymphocytes). Cytotoxic assay of test compounds was carried out by an adoption of the method described by Iqbal *et al* [22]. PBS was passed through 0.2 µm filter under laminar flow hood (kept sterile conditions) and then equal volumes of PBS and blood were taken in a sterile tube. Ficol solution (volume ratio 1:2) was carefully added at 165° angle to the mixture of PBS and blood. The tube was centrifuged at 2000 rpm at 4 °C for 30 minutes. The lower transparent portion was punctured with a syringe, and the liquid carefully removed and added to 5 ml of RPMI-1640 medium. The number of lymphocytes was counted in a haemocytometer under upright microscope. In the next step, 100 µl of the lymphocyte media was put into each well of a 96-well culture plate. The test compounds were added at doses of 100, 50, 25 and 10 µg/ml in DMSO, each in a final volume of 3 ml. Amphotericin-B (25 µg/ml, positive control) was used as reference drug, while negative control was *L. tropica* KWH23 promastigotes. Using a multipipette, 10 µl of promastigotes (containing  $1.4 \times 10^6$  promastigotes/ml) was added to 12 wells of the culture plate and placed in an incubator at 26 °C for at least 48 h. Haemocytometer was used to count viable lymphocytes and promastigotes under light microscope (lens 40x) at 24 and 48 hours. The cytotoxic tests were done in triplicate and the IC<sub>50</sub> for each compound was calculated [23].

### Statistical analysis

Results of *in vivo* anti-leishmanial assay of plant extract were expressed as mean % inhibition of parasite growth ± SD (n = 3). Cytotoxicity values were expressed as 50 % Inhibitory concentration (IC<sub>50</sub>) and analysed by non-linear regression analysis. For *in vivo* assays, mean lesion size (mm) and percentage cure were analysed by GraphPad Prism 5 software (GraphPad software, San Diego, CA) at 95 % confidence limit.

## RESULTS

Results of *in vivo* anti-leishmanial activities of compounds **1 - 5** analyzed in albino mice infected with 0.02 ml of clinically isolated *L. tropica* KWH23 having  $1.4 \times 10^6$  promastigotes, via intraperitoneal route, are shown Table 1. Mean lesion size decreased significantly from  $0.29 \pm 0.21$  mm to  $0.10 \pm 0.01$  mm after treatment with test compounds but that of the negative group reached  $1.5 \pm 0.02$  mm, whereas

it decreased in the Amphotericin (standard drug) group from  $0.85 \pm 0.05$  mm to  $0.17 \pm 0.01$  mm after 120 days. Luteolin showed the strongest anti-leishmanial activity. The mice groups that received apigenin, luteolin-4'-O- $\beta$ -D-glucopyranoside, and apigenin 4-O- $\beta$ -D-glucopyranoside had mean lesion sizes of  $0.29 \pm 0.21$  mm,  $0.25 \pm 0.09$  mm and  $0.21 \pm 0.03$  mm respectively, while corresponding percentage cure were 75.02, 78.51 and 80.19 % respectively. With Luteolin and lalioside mean lesion sizes were decreased to  $0.10 \pm 0.01$  and  $0.19 \pm 0.03$  mm, corresponding to 97.02 and 85.27 % cure, respectively.

### Cytotoxic effect

Results of cytotoxic activities of compounds 1-5 are summarized in Table 2. Compounds 3, 4 and 5 showed highest cytotoxic activities 10.27, 9.50 and 8.30  $\mu$ g/ml respectively. Compound 3 exhibited the greatest cytotoxicity against

lymphocytes, with IC<sub>50</sub> value of 10.27  $\mu$ g/ml (95 % C.I = 6.215-15.38). Compound 1 gave the lowest cytotoxic activity (IC<sub>50</sub> = 4.15  $\mu$ g/ml; 95 % C.I = 2.287-5.941), whereas compound 2 showed IC<sub>50</sub> value of 5.02  $\mu$ g/ml (95 % C.I = 2.136-7.087). In comparison, amphotericin-B (standard drug) displayed IC<sub>50</sub> of 1.072  $\mu$ g/ml (95 % C.I = 0.823-2.358).

### DISCUSSION

Compounds 1-5 were selected due to their promising *in vitro* anti-leishmanial effects [22] against *L. tropica*. All tested compounds showed a cure rate between 97.02 and 75.02 % at 3 mg/kg concentration (for 5 days) whereas cytotoxic effects against lymphocytes in terms of IC<sub>50</sub> values ranging between 4.15 and 10.27  $\mu$ g/ml. All the mice were cured at or after 120<sup>th</sup> day of infection, when compounds 1 - 5 were given;

**Table 1:** *In vivo* antileishmanial activity of plant phenolic compounds 1-5. Data represent mean lesion size (mm) with percentage cure rate with 95 % confidence intervals

Test compound	Dosing regimen <sup>a,b</sup> (For 5 Days) (mg/kg)	Variation of mean lesion size (mm)				Cure rate (with 95% Confidence intervals) <sup>c, d</sup>
		30 <sup>e</sup>	60 <sup>e</sup>	90 <sup>e</sup>	120 <sup>e</sup>	
Luteolin	3.0	$0.82 \pm 0.12$	$0.50 \pm 0.01$	$0.21 \pm 0.01$	$0.10 \pm 0.01$	97.022 (96.136-98.097)
Lalioside	3.0	$0.81 \pm 0.11$	$0.4 \pm 0.07$	$0.25 \pm 0.02$	$0.19 \pm 0.03$	85.27 (84.205-86.84)
Luteolin-4'-O- $\beta$ -D-glucopyranoside	3.0	$0.83 \pm 0.01$	$0.6 \pm 0.09$	$0.30 \pm 0.01$	$0.25 \pm 0.09$	78.51 (75.61-85.89)
Apigenin 4-O- $\beta$ -D-glucopyranoside	3.0	$0.84 \pm 0.10$	$0.71 \pm 0.03$	$0.35 \pm 0.03$	$0.21 \pm 0.03$	80.194 (77.002-87.671)
Apigenin	3.0	$0.82 \pm 0.01$	$0.60 \pm 0.02$	$0.40 \pm 0.01$	$0.29 \pm 0.21$	75.02 (74.96-76.39)
Amphotericin-B	15	$0.85 \pm 0.05$	$0.55 \pm 0.06$	$0.24 \pm 0.02$	$0.17 \pm 0.01$	95.00 (94.583-96.02)
Negative Control	3.0	$0.83 \pm 0.01$	$1.1 \pm 0.03$	$1.36 \pm 0.05$	$1.5 \pm 0.02$	0.000

a: Total dose 15 mg/kg; b: Route of administration: intraperitoneally (i.p); c: No: of mice cured/No: of mice Infected 6/6 (negative control 0/6); d: Mean survival time (Days)  $\geq$  60 (negative control  $\geq$  30); e: days post-infection

**Table 2:** Cytotoxic effect of plant phenolic compounds 1-5 against lymphocyte. Data represent IC<sub>50</sub> ( $\mu$ g/mL) with 95 % confidence intervals

Test compound	Lymphocyte cytotoxicity IC <sub>50</sub> ( $\mu$ g/mL) 95% confidence intervals
Luteolin	4.15 (2.287-5.941)
Lalioside	5.02 (2.136-7.087)
Luteolin-4'-O- $\beta$ -D-glucopyranoside	10.27 (6.215-15.38)
Apigenin 4-O-beta-D-glucopyranoside	9.50 (5.32-14.87)
Apigenin	8.30 (5.812-14.32)
Amphotericin-B	1.72 (0.823-2.358)

whereas after 60<sup>th</sup> day, mean lesion size was decreased up to  $0.4 \pm 0.07$  mm in mice. Luteolin

achieved a higher cure rate and decrease in lesion size than amphotericin B. Interspecies

variation in sensitivity against luteolin have been observed, e.g., it showed *in vitro* IC<sub>50</sub> of 12.5 µM against *L. donovani* [27,28] and reduced splenic parasitic load up to 80 % when 3.5 mg/kg (twice a week for 1 month) was given to *L. donovani* infected hamsters [29].

Luteolin decreased cell viability of human hepatoma HepG2 cells, result in decrease in cytotoxic activity up to 41 % [24]. In this study, lalioside, apigenin, luteolin-4'-O-β-D-glucopyranoside, and apigenin 4-O-β-D-glucopyranoside showed significant anti-leishmanial properties. Similar results have been previously reported. Tasdemir reported that apigenin showed *in vitro* anti-leishmanial activity against *L. donovani* with IC<sub>50</sub> values 1.9 µg/ml [28]. Apigenin showed cytotoxic activity at 8.30 µg/ml and induced hepatotoxicity at 100 and 200 mg/kg whereas it was non-toxic in 25 and 50 mg/kg [25]. In another study, apigenin decreased viability of human hepatoma HepG2 cells up to 42 % [24]. Apigenin 4-O-β-D-glucopyranoside showed cytotoxic activity at a concentration of 9.50 µg/ml and exhibited antitumor activity towards Hep G2, Hep 3B, MCF-7, A549 and MDA-MB-231 [26]. In the present study, lalioside and luteolin-4'-O-β-D-glucopyranoside showed greater cytotoxicity against lymphocytes, with IC<sub>50</sub> values of 5.02 µg/ml (95 % CL = 2.136-7.087) and 10.27 µg/ml (95 % CI = 6.215-15.38) respectively, compared with amphotericin B (IC<sub>50</sub> values = 1.072 µg/ml ;95 % CL = 0.823-2.358). The cytotoxic effects of these compounds may be due to the presence of phenolic hydroxyl groups [28]. Hydroxyl groups have affinity for proteins and may result in inhibition of microbial enzymes and plant NADH dehydrogenase [30,31]. It is evident that luteolin and luteolin-4'-O-β-D-glucopyranoside merit further investigations as suitable drug candidates against *L. tropica*.

Although different activities pertaining to the compounds studied are available in literature, the current study is the first report on the anti-leishmanial activities of compounds 2 – 4 and the first comprehensive *in vivo* study of cytotoxic activities of compounds 1-5 against *L. tropica* and mammalian cells.

## CONCLUSION

Luteolin is a potent anti-leishmanial agent, but is cytotoxic against lymphocytes. Luteolin-4'-O-β-D-glucopyranoside possesses significant anti-leishmanial activity, and is least toxic against lymphocytes. Lalioside exert anti-leishmanial activity while apigenin 4-O-β-D-glucopyranoside and apigenin have a moderate inhibitory effect

on intracellular amastigotes of *L. tropica* strain. Further studies on luteolin, luteolin-4'-O-β-D-glucopyranoside and other isolated compounds are necessary to investigate their mechanisms of action, specificity and structure-activity relationships.

## DECLARATIONS

### Acknowledgement

The authors would like to thank Prof Dr Fazal Subhan of Department of Pharmacy, University of Peshawar, Peshawar, Pakistan for providing the albino mice and other technical support.

### Conflict of Interest

No conflict of interest associated with this work.

### Contribution of Authors

The authors declare that this work was done by the authors named in this article and all liabilities pertaining to claims relating to the content of this article will be borne by them.

### Open Access

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

## REFERENCES

1. Alvar J, Ve'lez ID, Bern C, Herrero M, Desjeux P, Cano J, Jannin J, Boer MD. Leishmaniasis Worldwide and Global Estimates of Its Incidence. *PLoS ONE* 2012; 7(5): e35671. [PubMed]
2. Markle WH, Makboul K. Cutaneous leishmaniasis Recognition and Treatment. *Am Fam Physician* 2004; 69: 455-460
3. Pearson RD, Sousa ADQ. Clinical spectrum of leishmaniasis. *Clin Infect Dis* 1996; 22 (1): 1-13.
4. Brooker S, Mohammed N, Adil K, Agha S, Reithinger R, Rowland M, Ali I, Kolaczinski J. Leishmaniasis in refugee and local Pakistani populations. *Emerg Infect Dis* 2004; 10(9): 1681-1684.
5. Rowland M, Munir A, Durrani N, Noyes H, Reyburn H. An outbreak of cutaneous leishmaniasis in an Afghan

- refugee settlement in north-west Pakistan. *Trans Roy Soc Trop Med Hyg* 1999; 93(2):133-136.
6. Semwal RB, Semwal DK, Combrinck S, Cartwright-Jones C, Viljoen, A. *Lawsonia Inermis L. (henna): Ethnobotanical, Phytochemical and pharmacological aspects.* *J Ethnopharmacol* 2014; 155: 80-103. DOI:10.1016/j.jep.2014.05.042
  7. Sutthivaiyakit S, Unganont S, Sutthivaiyakit P, Suksamrarn A. *Diterpenylated and prenylated flavonoids from Macaranga denticulata.* *Tetrahedron* 2002; 58; 3619-3622. DOI:10.1016/S0040-4020(02)00296-X
  8. Baba K, Taniguchi M, Nakata K. *Studies on Angelica keiskei "Ashitaba".* *Foods Food Ingred J Japan* 1998; 178: 52-60.
  9. Fontenelle RO, Marais SM, Brito EH, Brilhante RS, Cordeiro RA, Lima YC, Brasil NV, Monteiro AJ, Sidrim JJ, Rocha MF. *Alkylphenol activity against candida spp. Microsporum canis: A focus on the antifungal activity of thymol, eugenol and O-methyl derivatives.* *Molecules* 2011; 16: 6422-6431.
  10. Cidade HM, Nascimento MS, Pinto MM, Kijjoa A, Silva AM, Herz W. *Artelastocarpin and carpelastofuran, two new flavones and cytotoxicities of prenyl flavonoids from artocarpus elasticus against three cancer cell lines.* *Planta med* 2001; 67 (9): 867-870.
  11. Meragelman KM, McKee TC, Boyd MR. *Anti-HIV prenylated flavonoids from Monotes africanus.* *J Nat Prod* 2001; 64(4): 546-548.
  12. Dufall KG, Ngadjui BT, Simeon KF, Abegaz BM, Croft KD. *Antioxidant activity of prenylated flavonoids from the West African medicinal plant Dorstenia mannii.* *J Ehtnopharmacol* 2003; 87(1): 67-72.
  13. Barron D, Balland C, Possety F, Ravanel P, Desfougeres A. *Prenyl flavonoids and renyl flavonoids and membrane-permeability,* *Acta botanica gallica* 1996; 143(6): 509-520.
  14. Mahmoud ZF, Abdel Salam NA, Khafagy SM. *Constituents of henna leaves (Lawsonia inermis L.) growing in Egypt.* *Fitoterapia* 1980; 51: 153-155.
  15. Liou JR, ElShazly M, Du YC, Tseng CN, Hwang TL, Chuang YL, Hsu YM, Hsieh PW, Wu CC, Chen SL, et al., *1,5-Diphenylpent-3-en-1-ynes and methyl naphthalene carboxylates from Lawsonia inermis and their anti-inflammatory activity.* *Phytochem* 2013; 88: 67-73. DOI:10.1016/j.phytochem.2012.11.010
  16. Afzal M, Al-Oriquat G, Al-Hassan JM. *Flavone glycosides from Lawsonia inermis.* *Heterocycles* 1980; 14: 1973-1976.
  17. Takeda Y, Fatope MO. *New phenolic glucosides from Lawsonia inermis.* *J Nat Prod* 1988; 51: 725-729.
  18. Hsouna AB, Trigui M, Culioli G, Blache Y, Jaoua S. *Antioxidant's constituents from Lawsonia Inermis leaves: Isolation, structure elucidation and antioxidative capacity.* *Food Chem* 2011; 125: 193-200. DOI: 10.1016/j.foodchem.2010.08.060
  19. Cuong NX, Nhiem NX, Thao NP, Nam NH, Dat NT, Anh HLT, Huong LM, Kiem PV, Minh CV, Won J-H, et al, *Inhibitors of osteoclastogenesis from Lawsonia inermis leaves.* *Bioorg Med Chem Lett* 2010; 20: 4782-4784.
  20. *National Research Council of The National Academy of Sciences. Guide for the Care and Use of Laboratory Animals: 8th edn. Washington, D.C.: The National Academies Press; 2010.*
  21. Jamal Q, Khan NH, Wahid S, Awan MM, Sutherland C, Shah A. *In-vitro sensitivity of Pakistani Leishmania tropica field isolate against buparvaquone in comparison to standard anti-leishmanial drugs.* *Exp Parasitol* 2015; 154: 93-97.
  22. Iqbal K, Iqbal J, Umair M, Farooq U, Iqbal MM, Qamar S, Bashir M. *Anti-leishmanial and cytotoxic activities of extracts from three Pakistani Plants.* *Trop J Pharm Res* 2016; 15(10): 2113-2119.
  23. Iqbal K, Iqbal J, Afreen MS. *Comparative Study on Antileishmanial and Cytotoxic activity of Lawsonia Inermis bark and Aloe Vera leaves.* *Int J Biol Pharm Allied Sci* 2016, 5(6), 1490-150.
  24. Lin JH, Lin YT, Huang YJ, Wen KC, Chen RM, Ueng TH, Liao CH. *Isolation and cytotoxicity of flavonoids from Daphnis Genkwae Flos.* *J Food Drug Anal* 2001; 9 (1): 6-11.
  25. Singh P, Mishra SK, Noel S, Sharma S, Rath SK. *Acute Exposure of Apigenin Induces Hepatotoxicity in Swiss Mice.* *PLoS ONE* 2012; 7(2): e31964. DOI:10.1371/journal.pone.0031964
  26. Lin AS, Chang FR, Wu CC, Liaw CC, Wu YC. *New cytotoxic flavonoids from Thelypteris torresiana.* *Planta Med* 2005; 71 (9): 867-870. DOI: 10.1055/s-2005-871292
  27. Mittra B, Saha A, Chowdhury AR, Pal C, Mandal S, Mukhopadhyay S, Bandyopadhyay S, Majumder HK. *Luteolin, an abundant dietary component is a potent anti-leishmanial agent that acts by inducing topoisomerase II-mediated kinetoplast DNA cleavage leading to apoptosis.* *Mol Med* 2000; 6: 527-541.
  28. Tasdemir D, Kaiser M, Brun R, Yardley V, Schmidt TJ, Tosun F, Rüedi P. *Antitrypanosomal and Antileishmanial Activities of Flavonoids and Their Analogues: In Vitro, In Vivo, Structure-Activity Relationship, and Quantitative Structure-Activity Relationship Studies.* *Antimicrob Agents Chemother* 2006; 50(4): 1352-1364.
  29. Salem MM, Werbovetz KA. *Natural Products from Plants as Drug Candidates and Lead Compounds against Leishmaniasis and Trypanosomiasis.* *Curr Med Chem* 2006; 13: 2571-2598.
  30. Prusky D, Keen NT. *Involvement of preformed antifungal compounds in the resistance of subtropical fruits to fungal decay.* *Plant Dis* 1993; 77: 114-119.
  31. Ravanel P, Creuzet S, Tissut M. *Inhibitory effect of hydroxyflavones on the exogenous NADH dehydrogenase of plant mitochondrial inner membranes.* *Phytochem* 1990; 29 (2): 441-445. DOI:10.1016/0031-9422(90)85093-U