

Research Article

Fast Dissolving Tablets of *Aloe Vera* Gel

Jyotsana Madan¹, AK Sharma², Ramnik Singh³

¹UP Technical University, Lucknow, India, ²MJP Rohilkhand, University, Bareilly, India, ³Sri Sai College of Pharmacy, Pathankot, India

Abstract

Purpose: The objective of this work was to prepare and evaluate fast dissolving tablets of the nutraceutical, freeze dried *Aloe vera* gel.

Methods: Fast dissolving tablets of the nutraceutical, freeze-dried *Aloe vera* gel, were prepared by dry granulation method. The tablets were evaluated for crushing strength, disintegration time, wetting time, friability, drug content and drug release. A 3² full factorial design was applied to investigate the combined effect of two formulation variables - amounts of microcrystalline cellulose and mannitol.

Results: The results of multiple regression analysis revealed that in order to obtain a fast dissolving tablet of the *Aloe vera* gel, an optimum concentration of mannitol and a higher content of microcrystalline cellulose should be used. A response surface plot was also provided to graphically represent the effect of the independent variables on the disintegration time and wetting time. The validity of the generated mathematical model was tested by preparing a check point batch.

Conclusion: This investigation has demonstrated that satisfactory fast dissolving *Aloe vera* gel tablets can be formulated. It also showed the potential of experimental design in understanding the effect of formulation variables on the quality of fast dissolving tablets.

Keywords: *Aloe vera*, Fast dissolving tablet, Factorial design, Mathematical model, Mannitol, Microcrystalline cellulose

Received: 06 August 2008

Revised accepted: 17 September 2008

*Corresponding author: Email: jmadan@hotmail.com, Mobile: +919463230415, Residence: +911862247868

INTRODUCTION

The genus, *Aloe*, belongs to the family, Liliaceae, and includes the species *Aloe barbadensis* Miller, commercially known as *Aloe vera*. *Aloe vera* has been used therapeutically for many centuries and is of particular interest due to its lengthy historic reputation as a curative agent and its widespread use in supplementary therapies. *Aloe vera* gel is the colorless gel contained in the inner parts of the fresh leaves¹. Chemical analysis has revealed that this clear gel contains amino acids, minerals, vitamins, enzymes, proteins, polysaccharides and biological stimulators. The *Aloe vera* gel, beginning in the 50's, has gained recognition as a base for nutritional drinks and foods²⁻⁴, as a moisturizer, and a healing agent in cosmetics⁵ and OTC drugs⁶.

Approximately one-third of the population, primarily, geriatric and pediatric populations, has swallowing difficulties, resulting in poor compliance with oral drug therapy⁷. Fast dissolving tablets offer the combined advantages of performance, convenience, rapid onset of action and patient compliance and allow administration of an oral solid dose form in the absence of water or fluid intake⁸. When placed on the tongue, it disintegrates instantaneously, releasing the drug which dissolves or disperses in the saliva⁹. They are prepared by techniques such as tablet molding, spray drying, lyophilization, sublimation, or addition of disintegrants¹⁰. Pharmaceutical formulators often face the challenge of finding the right combination of formulation variables that will produce a product with optimum properties. This study was undertaken to formulate a suitable fast dissolving nutraceutical tablet of freeze dried *aloe vera* gel (AVG), utilizing factorial design.

MATERIALS AND METHODS

Plant material

A. vera plants were collected (March 2003) and authenticated by Dr. C.S. Pandey of

Medicinal Plant Research and Development Centre, Govind Pant University of Agriculture and Technology, Pantnagar (Uttarakhand), India. A voucher specimen (AV-8) was retained in our museum for future reference.

Other materials

Croscarmellose sodium (CCS), crospovidone (CLP) and sodium starch glycolate (SSG) were purchased from S.D. Fine Chem., Mumbai, India. Microcrystalline cellulose - Avicel PH101 - hereinafter referred to as MCC, was procured from FMC Corporation, Philadelphia, USA. Mannitol was purchased from Merck India Ltd, Mumbai, India. Anhydrous lactose, talc, magnesium stearate and hydrochloric acid were obtained from CDH Chemicals, Delhi, India. Congo red reagent and methylene blue were acquired from Nice Chemicals Pvt. Ltd, Cochin, India.

Preparation of freeze-dried Aloe vera gel (AVG)

The inner mucilaginous parenchymatous tissues of leaves of *Aloe vera* plants were separated out with the help of a sterile knife. and homogenized in a blender (National blender, Matsushita Co. Japan) at 30 rpm. The homogenized mass was separated with a G3 sintered glass filter under vacuum, freeze-dried using a bench-top freeze-dryer (MC 2L, Cyberlab, USA) and subsequently stored at 4°C¹¹. The ratio of AVG to lyophilized powder was 200:1.

Preparation of AVG tablets

A preliminary screening of the disintegrants - croscarmellose sodium (CCS), crospovidone (CLP), sodium starch glycolate (SSG) and microcrystalline cellulose (MCC) - was conducted. Mannitol was incorporated as a soluble filler to improve palatability, impart a cooling sensation and sweet taste upon dissolution. Granulation was carried out by the dry granulation technique. All the ingredients were compressed and slugs of 0.8 g were produced at a compression force of 22.0 ± 1.0

kN using flat faced tooling 17 mm in diameter on a single punch tablet machine (Cadmach Machinery Ltd., Ahmedabad, India).

The slugs were then milled and the resulting granules sieved through sieve no. 20 USP. The granules were further mixed with a glidant-lubricant blend containing magnesium stearate(1%w/w) and talc (2% w/w) . The granules were compressed using a single punch tablet machine (Cadmach Machinery Ltd., Ahmedabad, India) fitted with 8 mm round standard concave punches. The tablet thickness was about 4.56 ± 0.06 mm.

Evaluation of the tablets

Hardness test

The crushing strength of the tablets (n=5) was measured using a Monsanto hardness tester (Sheetal Scientific industries, Mumbai , India).

Friability test

The friability of a sample of 20 tablets was measured using a Roche Friabilator (Electrolab, India). Twenty preweighed tablets were rotated at 25 rpm for 4 min. The tablet were then reweighed after removal of fines and the percentage of weight loss was calculated.

Wetting time

The wetting time of the tablets (n=6) was measured using a modified procedure described by Gohel et al¹². Five circular tissue papers (Dexina tissues, Gujarat, India) of 10 cm diameter were placed in a Petri dish (internal diameter 10 cm). Water (10 mL) containing methylene blue (10 % w/v), a water soluble dye, was added to the Petri dish. A tablet was carefully placed in the centre of the Petri dish and the time taken for the water to reach the upper surface of the tablets was noted as wetting time.

Disintegration time

This test was performed on 6 tablets. For disintegration time, one tablet was placed in the centre of the Petri dish (internal diameter 10 cm) containing 10 ml of water and the time taken by the tablet to disintegrate completely was noted¹².

Drug content uniformity

Colorimetric measurement of glucomannan in the AVG was used for determining drug content uniformity¹³. For the drug content, 10 tablets were weighed and triturated. A tablet triturate, equivalent to 2 mg of AVG, was weighed accurately and dissolved in 100 ml of distilled water and filtered. From this solution, 0.4 ml was transferred to a 10 ml test tube. To this, 4 ml of Congo red reagent (0.01 %) was added with mild vortexing. The mixture was left at room temperature for 20 min and absorbance was measured at 540 nm wavelength using UV-VIS spectrophotometer (SL-196, Elico Ltd). The amount of glucomannan was calculated by interpolating from the standard curve ($r^2 = 0.9747$).

Dissolution studies

The *in vitro* dissolution study was carried out in a USP dissolution test apparatus (TDT-OP Electrolab, Mumbai India), type 2 (paddle) with a dissolution media of 900 mL of 0.1 M hydrochloric acid at 50 rpm ($37 \text{ }^\circ\text{C} \pm 0.5 \text{ }^\circ\text{C}$). Samples (n=6) were withdrawn at the end of 30 min and the dissolution of drug was expressed as percent drug dissolved at the end of 30 min¹⁴.

Full factorial design

A 3² randomised factorial design was adopted to optimize the variables. In this design, two factors were evaluated each at 3 levels and experimental trials were performed at all 9 possible combinations¹⁵. The amount of disintegrant (MCC) and the soluble filler (mannitol) were chosen as independent variables. The disintegration time and wetting

time were selected as the dependent variables. The formulation and evaluation of factorial batches (F1 to F9) is shown in Table 2. The following statistical model incorporating interactive and polynomial terms was used to evaluate the responses:

$$Y = b_0 + b_1X_1 + b_2X_2 + b_{12}X_1X_2 + b_{11}X_1^2 + b_{22}X_2^2 \dots (1)$$

where Y is the dependent variable, b_0 is the arithmetic mean response of the 9 runs, and b_1 and b_2 are the estimated coefficients for the factors X_1 and X_2 respectively.

The main effects (X_1 and X_2) represent the average result of changing 1 factor at a time from its low to high value. The interaction terms ($X_1 X_2$) show how the response changes when 2 factors are simultaneously changed. The polynomial terms (X_1^2 and X_2^2) are included to investigate nonlinearity. The multiple regression analysis was performed followed by ANOVA to identify insignificant variables.

RESULTS

Hardness of the tablets was in the acceptable range of 2.56 to 3.55 kg/cm². Friability was 0.51 to 0.82 %. Drug content and release were 100±5% and 82.6-88.4 %, respectively. The results of the preliminary studies (Table 1) revealed that the tablets containing microcrystalline cellulose (MCC) exhibited rapid disintegration and wetting followed by tablets containing croscarmellose sodium (CCS), crospovidone (CLP), sodium starch glycolate (SSG) in that order.

Factorial design

The amount of disintegrant (MCC, X_1) and the soluble filler (mannitol, X_2) were chosen as independent variables in a 3² full factorial design. The disintegration time and wetting time were selected as the dependent variables. The data (Table 2) clearly indicates that disintegration time and wetting time are strongly dependent on the selected independent variables. The fitted equations (full and reduced) relating the responses, disintegration time and wetting time, to the transformed factor are shown in Table 3.

DISCUSSION

Preliminary trials

In order to select the best disintegrant, four disintegrants were studied in preliminary trials. The efficiency of disintegrants can be affected in varying magnitudes by the presence of a soluble filler in the tablet formulations¹⁶. This is expected due to the fact that the quantity of water penetrating into the tablet bed is limited. The soluble filler (mannitol) has good aqueous solubility, negative heat of solution and good wetting properties¹⁷. Mannitol will consume water partially, leaving only a part of the total water to penetrate into the tablet for the development of force necessary for disintegration. The results of the preliminary studies (Table 1) revealed that the tablets containing microcrystalline cellulose (MCC) exhibited rapid disintegration and wetting. MCC has good wicking and absorbing capacities¹⁸. Tablets of MCC disintegrated rapidly due to the rapid passage of water into the tablets resulting in the instantaneous rupture of the hydrogen bonds¹⁹. The delayed disintegration and wetting time of the tablets formulated using other disintegrants could be attributed to their slow water uptake and high gelling tendency. Based on the results of the preliminary study, MCC was selected as the disintegrant for further studies. The optimum concentration of MCC may be less than 12 %.

Factorial design

The polynomial equations can be used to draw conclusions after considering the magnitude of coefficient and the mathematical sign it carries (i.e., positive or negative). Table 4 shows the results of the analysis of variance (ANOVA), which was performed to identify insignificant factors. The high values of correlation coefficient for disintegration time and wetting time (Table 4) indicate a good fit. The equations may be used to obtain estimates of the response as a small error of variance was noticed in the replicates²⁰.

Table 1: Tablet formulation and evaluation of preliminary trials*

Ingredient	Batch code			
	A1	A2	A3	A4
Freeze dried aloe vera gel (mg)	150	150	150	150
Microcrystalline cellulose (mg)	25	-	-	-
Croscarmellose sodium (mg)	-	25	-	-
Sodium starch glycolate (mg)	-	-	25	-
Crospovidone (mg)	-	-	-	25
Mannitol (mg)	150	150	150	150
Anhydrous lactose q.s to---(mg)	400	400	400	400
Disintegration time (sec)	36.5	41.8	49.2	45.6
Wetting time (sec)	32.5	36.8	43.6	39.4
Hardness (Kg/cm ²)	3.55	3.24	2.56	2.96
Friability (% loss)	0.56	0.76	0.82	0.83
Drug content(%)	99.5	100.4	98.3	96.7
(%) Release	88.4	87.3	82.6	85.3

*All batches contained 2%w/w talc and 1%w/w magnesium stearate.

Table 2: 3² Full factorial design layout*

Batch code	Variable levels in coded form		Disintegration time	Wetting
	X ₁ (mg)	X ₂ (mg)	± SD (sec)	±SD(sec)
F1	-1	+1	36.75 ± 2.25	34.20 ± 0.75
F2	0	+1	36.33 ± 1.85	32.40 ± 1.08
F3	+1	+1	35.45 ± 3.32	32.66 ± 1.87
F4	-1	0	34.55 ± 3.12	30.54 ± 0.63
F5	0	0	34.00 ± 1.33	29.89 ± 1.23
F6	+1	0	33.24 ± 1.53	29.62 ± 0.89
F7	-1	-1	34.23 ± 2.45	31.66 ± 0.66
F8	0	-1	34.40 ± 4.23	30.90 ± 0.76
F9	+1	-1	34.14 ± 1.88	30.00 ± 1.23
Check point	+0.8	-0.2	33.6	29.2
Actual values				
Coded values	X ₁		X ₂	
-1	10		100	
0	20		150	
1	30		200	

*All batches contained 150 mg AVG, 2%w/w talc, 1%w/w magnesium stearate.

X₁ indicates amount of MCC(mg); X₂ is amount of mannitol (mg); SD = standard deviation.

Full and reduced models

The significance level of coefficients, b₁₂ and b₁₁, were found to be more than 0.05, hence they were omitted from the full model to generate the reduced model. The results of statistical analysis are shown in Table 3. The coefficients, b₁, b₂ and b₂₂, were found to be

significant at P < 0.05; hence they were retained in the reduced model. The reduced model was tested in portions to determine whether the coefficient b₁₂ and b₁₁ contributed significant information for the prediction of both disintegration time and wetting time. The results for the test of the model in portions are shown in Table 4. F-Statistics of the results of

Table 3: Summary of results of regression analysis*

For disintegration time						
Response (disintegration time)	b ₀	b ₁	b ₂	b ₁₂	b ₁₁	b ₂₂
FM	34.052	-0.450	0.960	-0.183	-0.303	1.287
RM	33.930	-0.450	0.960	-	-	1.287
For wetting time						
Response (Wetting time)	b ₀	b ₁	b ₂	b ₁₂	b ₁₁	b ₂₂
FM	29.761	-0.687	1.117	0.383	0.0300	1.953
RM	30.017	-0.687	1.117	-	-	1.953

*FM indicates full model; and RM, reduced model

Table 4: Calculations for testing the model in portions*

For disintegration time						
	DF	SS	MS	F	R ²	<i>F</i> _{calc} = 4.920 <i>F</i> _{table} = 9.55 DF = (2,3)
Regression						
FM	5	10.489	2.098	47.574	0.988	
RM	3	10.056	3.352	29.635	0.947	
Error						
FM	3	0.132	0.0441			
RM	5	0.566	0.113			
For wetting time						
	DF	SS	MS	F	R ²	<i>F</i> _{calc} = 0.198 <i>F</i> _{table} = 9.55 DF = (2,3)
Regression						
FM	5	18.239	3.648	18.397	0.968	
RM	3	17.942	5.981	33.511	0.953	
Error						
FM	3	0.595	0.198			
RM	5	0.892	0.178			

*DF indicates: degrees of freedom; SS, sum of squares; MS, mean of squares; F, Fischer's ratio; R², regression coefficient; FM, full model; and RM, reduced model.

ANOVA of full and reduced model confirmed omission of non-significant terms of equation 1. Since the calculated value (4.92 and 0.198) is less than the tabulated value (9.55, $\alpha=0.05$, df 2,3), it may be concluded that the interaction terms b₁₂ and the polynomial term b₁₁ do not contribute significantly to the prediction of both disintegration time and wetting time and, therefore, can be omitted from the full model. The results of multiple linear regression analysis (reduced model)

show that for both disintegration time and wetting time, the amount of MCC (X₁) had a negative effect while the concentration of mannitol (X₂) had a positive effect. It means that as the amount of MCC is increased, both the disintegration time and wetting time decreases, while as the amount of mannitol is increased both the disintegration time and wetting time would increase. Therefore, a high level of MCC and low level of mannitol should be selected for the rapid disintegration of the

tablets. The relationship between the dependent and independent variables was further elucidated using surface response plots. The data of the response surface plot (Figures 1 and 2)) demonstrate that both X_1 and X_2 affect the disintegration and wetting times. Based on the foregoing, discussion batch F6 (disintegration time: 33.24 sec and wetting time: 29.62 sec) was selected as a promising batch. A checkpoint batch was prepared at $X_1 = +0.8$ level and $X_2 = -0.2$ level. The values of disintegration and wetting times expected from the reduced model for this batch are 33.78 and 29.32 sec, respectively. Table 2 indicates the results were as expected. Thus, the statistical model is mathematically valid.




Figure 1: Response surface plot for disintegration time




Figure 2: Response surface plot for wetting time

CONCLUSION

The results of a 3^2 full factorial design revealed that the amount of microcrystalline cellulose and mannitol significantly affect the dependent variables - disintegration and wetting times. The present study has revealed the feasibility of optimization procedure in developing AVG fast dissolving tablets.

REFERENCES

1. Grindlay D, Reynolds T. *The Aloe vera phenomenon: A review of the properties and modern uses of the leaf parenchyma gel.* *J Ethnopharmacol*, 1986; 116:117-151.
2. Eshun K, He Q. *Aloe vera: a valuable ingredient for the food, pharmaceutical and cosmetic industries - a review.* *Crit Rev Food Sci Nutr*, 2004; 44: 91-96.
3. Atherton P. *Aloe vera: magic or medicine?.* *Nurs Stand*, 1998; 41: 49-54.
4. Vogler B K, Ernst E. *Aloe vera: a systematic review of its clinical effectiveness.* *Br J Gen Pract.* 1999; 49: 823-828.
5. Reynolds T, Dweck AC. *Aloe vera leaf gel: a review update.* *J Ethnopharmacol*, 1999; 68: 3-37.
6. Shelton RM. *Aloe vera. Its chemical and therapeutic properties.* *Int J Dermatol*, 1991; 30: 679-683.
7. Seager H. *Drug delivery products and the zydys fast dissolving dosage forms.* *J Pharm Pharmacol*, 1998; 50: 375-382.
8. Reddy LH, Ghosh B, Rajneesh. *Fast dissolving drug delivery systems: a review of the literature.* *Indian J Pharm Sci*, 2002; 64: 331-336.
9. Biradar SS, Bhagavati S T, Kuppasad I J. *Fast dissolving drug delivery systems: A brief overview.* *Internet J Pharmacol*, 2006; 4(2).
10. Chang R, Guo X, Burnside B, Couch R. *A review of fast dissolving tablets.* *Pharm Technol*, 2000; 24: 52-58.
11. Turner C E, Williamson DA, Stroud PA, Talley D J. *Evaluation and comparison of commercially available Aloe vera L. products using size exclusion chromatography with refractive index and multi-angle laser light scattering detection.* *Int Immunopharmacol*, 2004; 4:1727-1737.
12. Gohel M, Patel M, Amin A, Agarwal R, Dave R, Bariya N. *Formulation design and optimization of mouth dissolve tablets of nimesulide using vacuum drying technique.* *AAPS. Pharm.Sci.Tech*, 2004; 5(3): E36.
13. Eberendu AR, Luta G, Edwards JA, McAnalley BH, Davis B, Rodriguez S, Henry CR. *Quantitative colorimetric analysis of aloe polysaccharides as a*

- measure of Aloe vera quality in commercial products. *J AOAC Int*, 2005 ; 88: 684-691.
14. Patel DM, Patel M M. Optimization of Fast dissolving Etoricoxib tablets prepared by sublimation technique. *Indian J Pharm Sci*, 2008 ; 70: 71-76.
 15. Bolton S. *Pharmaceutical Statistics*. Ed 3, New York; Marcel Decker Inc. 1990, pp 234-236.
 16. Bolhuis GK, Zuurman K, Te Wierik GHP. Improvement of dissolution of poorly soluble drugs by solid deposition on a super disintegrant. II. The choice of super disintegrants and effect of granulation. *Eur J Pharm Sci*, 1997; 5: 63-69.
 17. Kizumi K, Watanabe Y, Morita K, Utoguchi N, Matsumoto M. New method of preparing high porosity saliva soluble compressed tablets using mannitol with camphor. *Int J Pharm*, 1997; 152 :127-131.
 18. Patel VF, Patel N M. Statistical evaluation of influence of viscosity of polymer and types of filler on dipyridamole release from floating matrix tablets. *Indian J Pharm Sci*, 2007; 69: 51-57.
 19. Jacob S, Shirwaikar AA, Joseph A, Srinivasan KK. Novel co-processed excipients of mannitol and microcrystalline cellulose for preparing fast dissolving tablets of glipizide. *Indian J Pharm Sci*, 2007; 69: 633-639.
 20. Sammour OA, Hammad MA, Megrab NA, Zidan AS. Formulation and optimization of mouth dissolve tablets containing Rofecoxib solid dispersion. *AAPST*, 2006;7: E55.